

Status of knowledge of the broad-nosed weevils of Colombia (Coleoptera, Curculionidae, Entiminae)

Jennifer C. Girón^{1,2}

¹ Department of Entomology, Purdue University, West Lafayette, IN, USA

² Natural Science Research Laboratory Museum of Texas Tech University, Lubbock, TX 79409, USA

Corresponding author: Jennifer C. Girón (entiminae@gmail.com)

Academic editor: P. Nunes-Silva | Received 16 October 2020 | Accepted 2 December 2020 | Published 30 December 2020

Citation: Girón JC (2020) Status of knowledge of the broad-nosed weevils of Colombia (Coleoptera, Curculionidae, Entiminae). Neotropical Biology and Conservation 15(4): 583–674. <https://doi.org/10.3897/neotropical.15.59713>

Abstract

Broad-nosed weevils in the subfamily Entiminae (Coleoptera: Curculionidae) are highly diverse, not only in terms of number of species, but also in their sizes, forms and colours. There are eight tribes, 50 genera and 224 entimine species recorded from Colombia: seven genera and 142 species are considered endemic and only a handful of species, which are recognised as pests of *Citrus* or potatoes, are broadly known. The large diversity of this subfamily in the country is only superficially known and even though genus level identifications are generally achievable, species identification remains quite challenging, due in part to limited access to broadly-scattered basic information. Summaries of available information and bibliographic resources for each of the tribes represented in Colombia are offered, along with a checklist of the species of Entiminae recorded from the country, obtained from literature and a pictorial key for tribal recognition. New combinations are proposed for eight species of the genus *Lanterius* Alonso-Zarazaga & Lyal. Information on the distribution of entimine species in Colombia is compiled for the first time, including complete references to each original description and available taxonomic revisions. About a third of the species of Entiminae remain as recorded from the country without specific locality information. In addition, genus level distributional maps are presented, generated from data obtained from four Colombian entomological collections. Lastly, some challenges for entimine identification in Colombia, which likely extend throughout the Neotropical region, are briefly discussed. This contribution aims, in part, to facilitate and promote entimine research in northern South America.

Resumen

Los gorgojos de pico corto en la subfamilia Entiminae (Coleoptera: Curculionidae) son altamente diversos, no sólo en términos del número de especies, sino además en tamaños, formas y colores. Existen ocho tribus, 50 géneros y 224 especies entiminos registradas en Colombia: siete géneros y 142 especies son consideradas endémicas, y sólo un puñado, las cuales son reconocidas como plagas de cítricos o papa, son ampliamente conocidas. La gran diversidad de esta subfamilia en el país sólo se conoce superficialmente y aunque identificar géneros es generalmente alcanzable, la identificación de especies sigue siendo un desafío, debido en parte al acceso limitado a información básica ampliamente dispersa. Se ofrecen resúmenes de información disponible y recursos bibliográficos para cada una de las tribus representadas en Colombia, junto con la lista de especies de Entiminae registradas para el país, obtenida a partir de la literatura, junto con una clave pictórica para el reconocimiento de tribus. Se proponen nuevas combinaciones para ocho especies del género *Lanterius* Alonso-Zarazaga & Lyal. Información sobre la distribución de las especies de Entiminae en Colombia se recopila por primera vez, incluyendo referencias completas a cada una de las descripciones originales y a las revisiones taxonómicas disponibles. Cerca de un tercio de las especies de Entiminae permanecen registradas para el país sin información específica de localidad. Adicionalmente, se presentan mapas de distribución a nivel de género, producidos a partir de datos obtenidos de cuatro colecciones entomológicas colombianas. Por último, algunas de las dificultades para la identificación de entiminos en Colombia, que probablemente se extienden a lo largo de la Región Neotropical, son brevemente discutidas. Esta contribución busca en parte facilitar y promover la investigación en Entiminae en el norte de Suramérica.

Keywords

Distribution, faunistics, literature, Neotropics, original description, South America, taxonomy

Palabras clave

Distribución, faunística, literatura, neotrópico, descripción original, Suramérica, taxonomía

Introduction

Weevils in the subfamily Entiminae Schönherr, 1823 (Schönherr 1823, col. 1138), commonly known as broad-nosed weevils (or “vaquitas” or “mulitas” in Spanish), constitute one of the most diverse, abundant and widespread groups of weevils worldwide, including well over 12000 described species (Oberprieler et al. 2007). The subfamily includes some of the most recognisable, charismatic and stunning weevil forms; for example, the genus *Eupholus* Boisduval, 1835 (Boisduval 1835, 363), from New Guinea, commonly known as “smurf-weevils” (see Van Dam et al. 2017), the genus *Pachyrhynchus* Germar, 1824 (Germar 1824, 336), in Southeast Asia (e.g. Háva and Rukmane 2018), sometimes called polka-dot weevils and the genera *Briarius* Fischer de Waldheim, 1829 (Fischer de Waldheim 1829, 88); formerly *Lamprocyphus* Marshall, 1922 (Marshall 1922b, 184; see Lanteri and del Río 2003) and *Entimus* Germar, 1817 (Germar 1817, 341), both from South America.

In taxonomic terms, the subfamily Entiminae has been recognised as such since Thompson (1992) and has been divided into as few as five tribes (Marvaldi 1997,

1998) to as many as 55 tribes (Alonso-Zarazaga and Lyal 1999); 54 according to Bouchard et al. (2011) is the most widely-accepted number of tribes. A summary of the taxonomic history of the Entiminae is provided by Marvaldi et al. (2014).

Biological generalities of entimines

Amongst curculionids, entimines are usually easy to recognise by the typical short and broad form of their rostrum, which is accompanied, in most cases, by conspicuous mandibular scars left by the breaking off of a deciduous process of the mandible at the time when the adult emerges from the pupal stage (Anderson and Howden 2002; see Fig. 1A). Entimines are phytophagous, trending to polyphagous and oligophagous; the larvae feed underground on the roots of plants, whereas the adults feed on the leaves (Marvaldi et al. 2014). Relatively few species of entimines (considering the whole diversity of the subfamily) are recognised as pests of economical importance; usually, a species only becomes a problem when populations are large. Especially relevant for Colombia are *Compsus viridivittatus* (Guérin-Ménéville, 1855) (Guérin-Ménéville 1855, 592), a *Citrus* pest in the tribe Eustylini (Cano et al. 2002a, b; Gallego et al. 2012, O'Brien and Peña 2012) and the 'Potato shooter' *Leuschenius vulcanorum* (Kirsch, 1889) (Kirsch 1889, 17) in the tribe Naupactini (Canchala 1992; Peña 2001; Cortázar Gómez et al. 2012).

A thorough summary of the natural history and morphology (of both adults and larvae) of the Entiminae is provided by Marvaldi et al. (2014). Additional information regarding morphological features of the adults and the larvae can be found in Marvaldi and Lanteri (2005) and Marvaldi (1997, 1998), respectively.

Most tribes are limited to particular biogeographic regions of the world (Marvaldi et al. 2014). Their distributions, in general, are more a reflection of habitat rather than host plant preferences (Anderson 1993, 2018; Marvaldi et al. 2014). The wide range of plants they feed on, paired with the strong association of species to particular habitats and geographic regions (e.g. Lanteri 1992; Anderson and Lanteri 2000; Franz and Girón 2009; Franz 2010b; Girón and Franz 2010; Franz 2011; Mazo-Vargas 2011, del Río and Lanteri 2011a; Girón and Franz 2012, del Río et al. 2015), makes entimines a potential model group for biogeographic studies and the identification of areas of endemism. There is also evidence of altitudinal stratification of species (Girón and Franz 2010).

As a consequence of their diversity and abundance, entimines are commonly found in the field, including urbanised areas and are, therefore, frequently found in large numbers in biological collections (see Girón and Cardona-Duque 2018). There is also a trend to find large numbers of individuals of single economically-important species in cultivars, whereas only few specimens of several different species can be found associated with forested or less disturbed areas (pers. obs.). Furthermore, it is common that specimens in collections are only identified to family or, at best, sorted to subfamily, as is often the case in Colombia (see Girón and Cardona-Duque 2018).




Figure 1. Select morphological features of Entiminae: **A, G** ventral view of rostrum: **A** right deciduous process still attached, left mandible with mandibular scar, maxillae fully covered by prementum (Naupactini) **G** mandibular scars well developed, maxillae partially covered by prementum (*Anypotactus* sp.) **B, C** frontal view of head indicating position of epistoma, nasal plate and mandibular scars: **B** Naupactini, with median furrow **C** Eustylini, with median fovea **D–F** lateral view of head and anterior section of prothorax: **D** Tanymericini, with anterior margin of prothorax straight and postocular setae clustered as a tuft **E** Eustylini, with anterior margin of prothorax slightly sinuate and without postocular setae **F** Lordopini, with anterior margin of prothorax strongly sinuate as to form well-developed postocular lobe, with postocular setae forming a fringe **H–K** profemora in frontal view (top end of drawings articulate to coxa, bottom end articulate to tibia): **H** *Compsus* sp., regularly shaped profemur **I** *Eustylus* sp., toothed profemur **J** *Hadromeropsis* sp. enlarged profemur **K** *Anypotactus* sp., clavate and toothed profemur **L, M** apical region of left metatibia (left margin on drawing is anterior (inner) in the beetle; right margin on drawing is posterior (outer) in the beetle), black triangles indicate posterior corner of metatibia: **L** Eustylini, posterior corner of metatibia rounded **M** Naupactini, posterior corner of metatibia angulate **N–P** anterior section of elytra in dorsal view: **N** shoulders absent **O** shoulders oblique, weakly developed **P** shoulders well developed.

Identification challenges

One of the most relevant references for Neotropical Broad-nosed weevils is the 'Biologia Centrali-Americana' by Sharp and Champion (1911), although it includes only scarce information on Colombian taxa. Even though many genera have been revised (see below under the treatment for each tribe), species identification of Neotropical and, especially northern South American, entimines remains highly challenging (e.g. Girón 2006). For Colombia, in particular, the fact that most type specimens are deposited in collections overseas, combined with the lack of identified material in national collections and compounded by a lack of specialists in the region, makes species identification a very difficult task at this time. In addition, one of the main difficulties for identifying entimine taxa is the limited access to information, which has been changing in the digital era. Many publications, including original descriptions and revisions are currently available online, some of them for free through the Biodiversity Heritage Library (<https://www.biodiversitylibrary.org/>). Nevertheless, some key publications with full references linking to those original descriptions (e.g. Wibmer and O'Brien 1986a; Alonso-Zarazaga and Lyal 1999) remain available only on paper and are essentially inaccessible to the broad community, especially in Latin American countries. An effort has been made in this paper to incorporate references and links to as many relevant publications as possible.

This contribution compiles and summarises the available information for the Entiminae recorded from Colombia. A general pictorial key to diagnose tribes is given. For each tribe represented in the country, a list of morphological characters for tribal recognition is offered, along with distributional information and bibliographic resources. A list of species is provided, including bibliographic records of distributional data within the country, where available. In addition, genus-level information has been recorded from specimens identified in Colombian national entomological collections, further providing information about distributions of entimines in the country.

Materials and methods

Species list

The list of species of Entiminae recorded from Colombia, as well as general species distributions, were extracted from the annotated checklist of the weevils of South America by Wibmer and O'Brien (1986a), the list of species of Entiminae by Morrone (1999), in addition to more recent revisions or new descriptions, where available. Tribal concepts and general distributions of genera are in accordance with Alonso-Zarazaga and Lyal (1999). The full list of species is included here as Suppl. material 1 and available as a checklist via GBIF (Girón 2020, <https://doi.org/10.15472/jdwfao>); this online resource will be updated whenever new species are added to the Colombian fauna of Entiminae.

Only presence in Colombia was obtained from the main references (Wibmer and O'Brien 1986a; Morrone 1999 and Alonso-Zarazaga and Lyal 1999); therefore, original descriptions for each species were checked to obtain distributional records within the country. Localities taken from the 1800s' to mid-1900s' publications might not reflect current departments or municipalities (e.g. 'Bogotá' as a locality in 1857 does not strictly correspond to current 'Cundinamarca: Bogotá'). In those cases, locality is quoted as presented in the original description. Department was added for unambiguous localities (e.g. Antioquia was added to original localities recorded as 'Medellin'; Cundinamarca added for Fusagasugá; Meta added for Villavicencio, Boyacá added for Muzo). Localities in square brackets '['] in the list of species indicate locality spelling in the original description. Type locality is recorded as 'Bogotá' for species described by Kirsch (1868) given the title of the paper: "Beiträge zur Käferfauna von Bogotá". A question mark '?' indicates that the locality record is doubtful and needs confirmation. Specific localities from countries other than Colombia are omitted here. For some of Schönherr's taxa, the page number corresponds to a column number in the publication and is indicated by 'col.' (e.g. Entiminae Schönherr 1823, col. 1138).

Material examined

Specimens from entomological collections in Colombia were revised and distributional data at the generic level were recorded (Suppl. material 2). The insect collection abbreviations are adopted, for the most part, from Evenhuis (2020) as follows:

IavH	Instituto Alexander von Humboldt, Villa de Leyva, Colombia, https://doi.org/10.15472/vmpedy [databased July 2013]
MEPB	Museo Entomológico de Piedras Blancas, Comfenalco Antioquia, Colombia, https://doi.org/10.15472/xblw2v [databased March 2013]
MPUJ	Museo Javeriano de Historia Natural Lorenzo Uribe, S.J., Bogotá, Colombia, https://doi.org/10.15472/xmukx8 [partly databased October 2015]
MUSENUV	Museo de Entomología de la Universidad del Valle, Cali, Colombia, https://www.gbif.org/publisher/85be57ed-f187-49c9-b7ff-eea622e06217 [databased December 2006]

A few additional records were obtained from ASUCOB (Arizona State University Charles W. O'Brien Collection, Arizona, USA <https://doi.org/10.15468/eeqtx0>). Information obtained from collections is summarised for each tribe and identified genus.

Genus-level identifications were obtained using available keys (van Emden 1944a, b) and diagnoses (Sharp and Champion 1911) and confirmed by consulting experts in Curculionidae (Dr. Robert Anderson, Canadian Museum of Natural History, Ottawa, Canada and Dr. Charles O'Brien, USA), who have access to reference collections. The recognition section presented here for each tribe applies to entimines seen in Colombian collections and might not reflect the entire variation across tribes and genera. In most cases, there is more than one morphospecies with-

in each genus at each collection. Species were not identified. There remain specimens recognised as entimines, but not identified to tribe. Relevant information for entimine tribes represented in Colombia was compiled and is presented here. A general pictorial key is provided here just as a way to narrow down identifications (Fig. 2). Preliminary identifications using this key should be checked against the lists of characters offered for each tribe for confirmation.

Figures

Line drawings were produced by tracing over photographs into drawing software. Figures are grouped to show morphological variation or to keep relevant tribal information together. Maps are based on data from collections. The record density map (Fig. 3) was created using R version 3.5.2 (R Core Team 2018), interfaced through RStudio version 1.1.463 (RStudio Team 2016) with the R package ‘tmap’ (Tennekes 2018). Distribution maps in Figs 3–10 were created using SimpleMappr (Shorthouse 2010); each map is accompanied by sketches of the general appearance of the head in dorsal view of some representative specimens of the tribe as seen in Colombian collections.

Results

There are eight tribes, 50 genera, and 224 species of Entiminae recorded from Colombia. Seven genera and 142 species are considered endemic to the country (Wibmer and O’Brien 1986a; Morrone 1999; Girón and Cardona-Duque 2018; see species list, Suppl. material 1 or Girón 2020 – <https://doi.org/10.15472/jdwfao>). A general pictorial key to recognise these eight tribes is presented in Fig. 2. The list of species with their distribution in Colombia is presented. The genera *Pandeleiteius* Schönherr, 1834 (Schönherr 1834, 129; Tanymecini, 44 spp.), *Compsus* Schönherr, 1823 (Schönherr 1823, col. 1140; Eustylini, 39 spp.), *Hypsonotus* Germar, 1824 (Germar 1824, 367; Lordopini, 17 spp.) and *Exorides* Pascoe, 1881: 43 (Pascoe 1881, 43; Eustylini, 14 spp.) are the most diverse in the country (numbers in parentheses indicate numbers of species recorded from Colombia).

Diversity by department

A total of 749 records of entimines were obtained from the revised collections (Suppl. material 2), with 248 (33%) from Valle del Cauca, followed by Magdalena (75, 10%), Antioquia (62, 8.3%), Risaralda (48, 6.4%), Boyacá (47, 6.3%), Nariño (35, 4.7%), Bolívar (31, 4.1%), Meta (27, 3.6%), Quindío (22, 3%), Cauca, Cundinamarca and Amazonas, each with 18 records (2.4%), followed by Tolima (11, 1.5%), Chocó (9, 1.2%), Santander (7, 1%), Putumayo and Vaupés with 6 each (0.8%), Norte de Santander (5, 0.7%), Vichada (3, 0.4%), Guaviare (2, 0.3) and Cesar with only one record (0.1%). The Departments of Atlántico, Caquetá, Córdoba, Huila, La Guajira, Sucre, Arauca, Guainía and San Andres, Providencia and Santa Catalina have no entimine occurrences from the studied collections (Fig. 2).




Figure 2. Pictorial key for the recognition of Tribes represented in Colombia. Preliminary identifications achieved using this key should be checked against the lists of characters offered for each tribe for confirmation.

Tribe Anypotactini Champion, 1911

Fig. 4

Recognition. Usually small (approx. 4–11 mm); scale coverage composed by appressed, mostly non-overlapping scales, usually uniform (in density and colour), usually brown (some species iridescent green or blue); usually with thick and erect




Figure 3. Record density map of Entiminae in Colombia. The colour gradient in the map represents the number of records by Department recorded from entomological collections (see Materials and Methods). Departments in white indicate no records and Departments increasingly darker with larger numbers of records (numbers in parentheses): Valle del Cauca (248), Magdalena (75), Antioquia (62), Risaralda (48), Boyacá (47), Nariño (35), Bolívar (31), Meta (27), Quindío (22), Cauca, Cundinamarca and Amazonas, (18 each), Tolima (11), Chocó (9), Santander (7), Putumayo and Vaupés (6 each), Norte de Santander (5), Vichada (3), Guaviare (2) and Cesar (1). Atlántico, Caquetá, Córdoba, Huila, La Guajira, Sucre, Arauca, Guainía and San Andres, Providencia and Santa Catalina have no entimine occurrences from the studied collections. The data used to create this map are available in Suppl. material 2.

scale-like setae, rather uniformly distributed all over the surface; head (including rostrum) subrectangular [e.g. some *Anypotactus* Schönherr, 1840 (Schönherr 1840b, 299; Fig. 4B) and *Polydacrys* Schönherr, 1834 (Schönherr 1834, 130; Fig. 4D)] to conical [*Phanasora* Pascoe, 1881 (Pascoe 1881, 38; Fig. 4C), *Prepodellus* Kirsch, 1868 (Kirsch 1868, 239) and some *Anypotactus*], usually longer than wide; surface of rostrum flat, depressed or undulate, often with longitudinal carinae (Fig. 4B–D); nasal plate well developed (see Fig. 1B, C), forming an elevated, large and smooth plate in *Polydacrys* (Fig. 4D); mentum not covering the maxillae completely (Fig. 1G; except in *Polydacrys*, see Fig. 1A); antennal scape slender, at rest passing well below eye, often reaching or surpassing anterior margin of prothorax; anterior margin of prothorax in lateral view straight to oblique, never forming postocular lobe, lacking postocular setae [except in *Hyphantus* Germar, 1824 (Germar 1824, 334)]; elytral shoulders and hindwings weakly- to well-developed (Fig. 1N–P); legs relatively slender, femora clavate, often toothed (Fig. 1K).

Anypotactines can be confused with some small Naupactini, from which they can be recognised by the shape of the apical region of the metatibia, which is rectangular and fringed by flat spines along ventral and posterior margins in Naupactini (Fig. 1M, rounded in Anypotactini, see Fig. 1L). The general appearance of some Anypotactini resembles members of Tanymericini (Fig. 1D, with postocular setae) and can be differentiated, most of the time, by the lack of postocular setae in Anypotactini.

Diversity. The tribe contains 11 extant genera with 81 species described, all distributed in the New World (Wibmer and O'Brien 1986a, 44, Alonso-Zarazaga and Lyal 1999, 145). Seven genera represented by 11 species have been recorded from Colombia. *Hypsometopus* Kirsch, 1868 (Kirsch 1868, 223), represented by a single species, is considered endemic to the country; seven additional anypotactine species are considered endemic as well.

General distribution. The tribe ranges from south-western USA to Argentina and Chile and some Caribbean islands, including the following countries: Argentina, Bolivia, Brazil, Chile, Colombia, Ecuador, Peru, Paraguay, Uruguay, Venezuela; Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama; Cuba, Dominica, Granada, Guadeloupe, Puerto Rico, Saint Vincent; SW USA (Texas) (Alonso-Zarazaga and Lyal 1999). Most genera are distributed in Central America and northern South America, but the largest genus, *Hyphantus* Germar, 1824 (Germar 1824, 334; 45 species; see Vaurie 1963), is distributed in southern Brazil, Argentina, Paraguay and Uruguay.

Distribution in Colombia. Within the revised collections, anypotactines were recorded from Bolívar, Boyacá, Caldas, Cundinamarca, Magdalena, Quindío, Risaralda, Santander and Valle del Cauca (Fig. 4A). *Anypotactus* and *Polydacrys* were the most commonly found anypotactine genera in the revised collections; other anypotactines were found but remain identified to tribe only (Suppl. material 2; Fig. 4A).

Bibliographic resources. The best resource available for the tribe as a whole is the *Biologia Centrali-Americana* (Sharp and Champion 1911, 215, as 'Anypotactina'). It includes a brief diagnosis of the tribe and diagnoses of the Central American




Figure 4. Distribution and some head variants of Anypotactini: **A** distribution map of records for *Anypotactus*, *Bothynodontes*, *Phanasora*, *Polydacrys*, *Prepodellus* and undetermined Anypotactini in Colombia **B–D** head in dorsal view: **B** *Anypotactus* **C** *Phanasora* **D** *Polydacrys*.

genera and species (with some keys). There is a key in van Emden (1944b, 506) to identify the tribe, as well as a key to genera (van Emden 1944b, 510). The only genus of the tribe that has been revised is *Hyphantus* (Vaurie 1963), but it is not represented in Colombia.

Natural history. Except for a few plant-association records for *Hyphantus* (Vaurie 1963) and records of an unidentified species of *Bothynodontes* on *Alnus jorulensis* (Betulaceae) in Caldas and *Baccharis* sp. (Asteraceae) in Tolima (Bustillo and Villegas Isaza 1986), the natural history of Anypotactini remains largely unknown.

Remarks. A large proportion of anypotactine records in Colombia are concentrated in the mountainous regions (Fig. 4A). Some specimens amongst the undetermined

Anypotactini from Caldas (Manizales, Río Blanco; MUSENUV) likely belong to an undescribed genus (Dr. Robert Anderson, pers. comm.). Specimens recorded from lowlands, especially from the north coast, share affinities with material seen from Central America at ASUCOB. One species of *Anypotactus* was included in a morphology-based phylogenetic analysis (Girón and Franz 2010), resulting as sister to *Apodrosus* Marshall, 1922 (Marshall 1922a, 59; Polydrusini, exclusively Caribbean); no anypotactines have been included in molecular-based phylogenetic analyses to date.

Tribe Entimini Schönherr, 1823

Fig. 5

Recognition. Medium to large (approx. 7–45 mm), often very robust; scale coverage either nearly uniform and dense or forming patches across surface, highly variable in colouration including brown, white, purple and iridescent tones of green and blue; head (including rostrum) subrectangular, usually longer than wide (Fig. 5B, C); rostrum usually apically broadened (Fig. 5C); anterior margin of prothorax in lateral view strongly sinuate forming conspicuous postocular lobe (Fig. 5D), often with fringe of postocular setae (Fig. 5D); elytral shoulders well developed, sometimes projected from general outline of body; hindwings well-developed; surface of elytra may be granulate or bear protuberances or tubercles.

Members of Entimini are generally easily distinguishable by their usually large size and the general bulky shape of their bodies [*Cydianerus* Schönherr, 1840 (Schönherr 1840a, 737), *Entimus*, *Polyteles* Germar, 1829 (Germar 1829, 358)] or the characteristic shape of their heads with narrow frons and rostrum with lateral margins apically diverging [*Cydianerus*, *Rhigus* Schönherr, 1823 (Schönherr 1823, col. 1138)]. A more detailed diagnosis is offered by Vaurie (1951).

Diversity. The tribe contains seven Neotropical genera with 46 species described to date, two of them fossil (Wibmer and O'Brien 1986a, 99, Gaiger 2001; Morrone 2002; Vanin and Gaiger 2005; Poinar et al. 2017). Three genera, represented by four species, have been recorded from Colombia, one of them fossil (Poinar et al. 2017).

General distribution. The tribe ranges from Mexico to Argentina, with no representatives in the Caribbean islands. Entimini can be found in the following countries: Argentina, Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Paraguay, Peru, Uruguay, Venezuela; Costa Rica, El Salvador, Honduras, Mexico, Nicaragua, and Panama (Alonso-Zarazaga and Lyal 1999). All genera are mainly represented in Brazil.

Distribution in Colombia. In Colombian collections, representatives of Entimini (genus *Rhigus*) have been recorded from Amazonas and Tolima (Girón and Cardona-Duque 2018, 181). The distribution in the country for the rest of Colombian Entimini has not been recorded in literature. There is one record of the genus *Cydianerus* from Sucre in iNaturalist (<https://www.inaturalist.org/observations/39591579>).




Figure 5. Distribution and some head variants of Entimini: **A** distribution map of records for *Rhigus* in Colombia **B, C** head in dorsal view: **B** *Entimus* **C** *Rhigus* **D** head and anterior section of prothorax in lateral view of *Rhigus* showing well-developed postocular lobe.

Bibliographic resources. Entimini is one of the better-studied groups of Neotropical broad-nosed weevils. Sharp and Champion (1911, 300), with emphasis on Central American fauna, offer some diagnostic features for a wider concept of Entimini (as 'Entimina') which includes some genera currently placed elsewhere. Bovie (1908a, 2) offered a key to genera including illustrations (Bovie 1908a, plate 9). Morrone (1999) treated 'Entimina' in a broader sense, including 'Entimina', 'Lordopina', 'Hypsonotina', 'Promecopina' and 'Eudiagogina' of Wibmer and O'Brien (1986a). Some genera of the tribe have been revised: *Entimus* (Vaurie 1951; Morrone 2002), *Rhigus* (Gaiger 2001) and *Phaedropus* Schönherr, 1823 (Schönherr 1823, col. 1140; Vanin 1983; not recorded from Colombia). A morphology-based phylogeny, key to the current genera and details on their morphology were presented by Vanin and Gaiger (2005).

Remarks. Even though members of Entimini are amongst the largest and most striking weevils in the subfamily, they are extremely rare in Colombian collections; only a few specimens of *Rhigus* have been found in the revised collections along with a few other unidentified Entimini. Romo and Morrone (2011) analysed biogeographic trends in the tribe. With the exception of the life cycle for *Entimus nobilis* (Olivier, 1790) (Olivier 1790, 525, Bruch 1932), the natural history of members of Entimini remains largely unknown. Plant associations have only been recorded for one species of *Phaedropus* (Vanin 1983) and one species of *Rhigus* (Gaiger 2001). There is one fossil species of *Cyodianerus* recently described from Colombia (Poinar et al. 2017) and one fossil species of *Entimus* described from the USA (Scudder 1876). There is a recent study on the morphological variation and sexual dimorphism in *Cyodianerus latruncularius* (Perty, 1832) (Perty 1832, 70, Regueira et al. 2020). *Rhigus speciosus* has been recently recorded from Colombia (Girón and Cardona-Duque 2018) and Peru (Gillett and Barr 2020). The type species for *Entimus*, *Entimus imperialis* Forster, 1771 (Forster 1771, 34), has been the focus of studies regarding the structural colouration of its scales which, as in other entimines, contain three-dimensional photonic crystals with diamond-type structure (e.g. Wilts et al. 2012). *Polyteles* Germar, 1829 (Germar 1829, 358) has not been recorded from Colombia, but its presence in the south of the country is likely, given that the genus is distributed in Argentina, Bolivia, Brazil, Ecuador, Paraguay, Peru and Uruguay.

Tribe Eudiagogini LeConte, 1874

Fig. 6

Recognition. Small weevils (approx. 3–8 mm); scale coverage usually light to dark brown, forming patterns on dorsal surface; iridescent areas with blue or green scales or reddish or yellowish stripes may be present in some genera; setae along surface may also be present; surface of body smooth and even; head (including rostrum) subrectangular, usually nearly as long as wide (Fig. 6B, C); rostrum robust, usually nearly parallel-sided (Fig. 6B, C); nasal plate (see Fig. 1B, C) usually well developed; anterior margin of prothorax in lateral view strongly sinuate forming conspicuous broad postocular lobe (Fig. 6D); elytral shoulders moderately to well-developed (e.g. Fig. 1O, P); legs relatively stout, femora never toothed [except in *Eurysaces* Schönherr, 1840 (Schönherr 1840b, 313), not present in Colombia].

Eudiagogines are easily recognisable by their small size, well-developed postocular lobes and overall robustness. They could potentially be confused with anypotactines because of their size, but their postocular lobe sets eudiagogines apart. They can resemble some small Entimini, but eudiagogines have a comparatively shorter and stouter rostrum. Additional characters to define the tribe can be found in Lacordaire (1863, 384, in French).

Diversity. The tribe contains ten extant genera with 100 species (including two fossil species in Dominican amber; Poinar and Brown 2011; Poinar and Legalov 2017)




Figure 6. Distribution and some head variants of Eudiagogini: **A** distribution map of records for *Promecops* and undetermined eudiagogines in Colombia **B**, **C** head in dorsal view: **B** *Promecops* **C** *Colecerus* **D** head and anterior section of prothorax in lateral view of *Promecops* showing well-developed postocular lobe.

described to date, all distributed in the New World (as 'Promecopini' in Wibmer and O'Brien 1986a, 81, O'Brien and Kovarik 2001). Two genera, *Colecerus* Schönherr, 1840 (Schönherr 1840a, 927) and *Promecops* Sahlberg, 1823 (Sahlberg 1823, 30), represented by six species, have been recorded from Colombia; four species are endemic.

General distribution. The tribe ranges from the USA to Argentina, including some Caribbean islands. Eudiagogini can be found in the following countries: Argentina, Bolivia, Brazil, Chile, Colombia, Ecuador, French Guiana, Guyana, Paraguay, Peru, Uruguay, Venezuela; Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama; Grenadines, Guadeloupe, St. Vincent; NC, NE, SE, SW USA (Alonso-Zarazaga and Lyal 1999).

Distribution in Colombia. Representatives of the tribe have been recorded in collections from Amazonas, Bolívar, Chocó, Magdalena, Meta, Risaralda, Tolima, Valle del Cauca, and Vaupés (Suppl. material 2, Fig. 6A).

Bibliographic resources. The original diagnosis for Eudiagogini (then named Promécopides) was given by Lacordaire (1863, 384, in French). In the *Biologia Centrali-Americana* (Sharp and Champion 1911, 300) eudiagogine genera were considered part of an extended concept of Entimini (as ‘Entimina’). Morrone (1999) treated ‘Eudiagogina’ and ‘Promecopina’ as part of ‘Entimina’. A revision of *Eudiagogus* Schönherr, 1840 (Schönherr 1840b, 307), was published by Warner (1979). LeConte and Horn (1876, 108) treated the genera recorded from North America. Voss (1934, 74, in German, as ‘Promecopina’), provided a key to genera (containing genera of Entimini) and keys to species for some eudiagogine genera.

Remarks. *Promecops* contains over 60% of the species and is also the most widespread genus of the tribe; some species are considered economically important in soybean cultivars in Argentina (Lázaro et al. 1997, 1998) and Brazil (Rocha Barreto and Cavalet 2016). One species of *Colecerus* has been associated with cultivated Pecan trees in Mexico (Soto-Hernández and Barros-Barrios 2018). In North America, *Eudiagogus* has been associated with legumes in the genus *Sesbania* (Warner 1979); *Eudiagogus* has not been recorded from Colombia, but its presence in the south of the country is likely, given that the genus is distributed in Argentina, Bolivia, Brazil, Ecuador, Paraguay, Peru, Uruguay, Costa Rica, Honduras, Mexico and USA. *Promecops* is the only eudiagogine genus recorded from the revised Colombian collections; other eudiagogines were located, but remain undetermined (Suppl. material 2, Fig. 6A).

Tribe Eustylini Lacordaire, 1863

Fig. 7

Recognition. Medium to large weevils (approx. 10–25 mm); scale coverage highly variable in presence, density and colouration; iridescent scales, erect setae or waxy secretions are frequently present; surface smooth and even or strongly sculptured and irregular; head (including rostrum; Fig. 7C–F) subrectangular, nearly as long or longer than wide; eyes small to mid-sized, slightly dorsally positioned; frons usually as wide as or narrower than interantennal distance, often bearing median fovea (see Fig. 1C); rostrum nearly parallel-sided or broadened apically; dorsal surface of rostrum with variable elevations or depressions, including longitudinal carinae or oblique fossae; antennal scrobe generally fully visible in dorsal view (Fig. 7C–F); nasal plate (see Fig. 1B, C) usually well developed, either depressed, flat or elevated regarding surface of rostrum; anterior margin of prothorax in lateral view straight, seldom slightly sinuate (Fig. 1E), never forming conspicuous postocular lobe; postocular setae may be present, if so, forming a fringe instead of a tuft (compare Fig. 1D vs. 1F); elytral shoulders usually well-developed, absent in *Brachyomus* La-




Figure 7. Distribution and some head variants of Eustylini: **A, B** distribution maps of Eustylini in Colombia: **A** records for *Brachyomus*, *Compsus*, *Eustylus* and *Exophthalmus* **B** records for *Exorides*, *Oxyderces*, *Synthlibonotus*, *Xestogaster* and undetermined eustylines **C–F** head in dorsal view: **C** *Exophthalmus* **D** *Synthlibonotus* **E** *Exorides* **F** *Eustylus* with characteristic broad antennal scape.

cordaire, 1863 (Lacordaire 1863, 130), reduced in some *Compsus* and *Exorides* Pascoe, 1881 (Pascoe 1881, 43) (see Fig. 1N–P); tubercles and apical projections may be present on elytra; femora usually not toothed (Fig. 1H; except in some *Eustylus* Schönherr, 1842 (Schönherr 1842, 40), Fig. 1I).

Eustylines are relatively easy to recognise amongst South American entiminae by their size, frequently iridescent/bright colourations and the shape of their head (Fig. 7C–F). They may be confused with similarly-coloured Naupactini, but the shape of the head, eyes, nasal plate and apex of metatibia [compare Fig. 1L (Eustylini) vs. 1M (Naupactini)] differentiate them.

Diversity. Eustylini contains 25 genera with 336 species described to date. Eight genera, represented by 73 species, have been recorded from Colombia, 52 of which are endemic.

General distribution. Eustylini ranges from south-western USA to Argentina, with its highest diversity in the Caribbean, Central America and northern South America, including the following countries: Argentina, Bolivia, Brazil, Colombia, Ecuador (incl. Galapagos Islands), French Guiana, Guyana, Paraguay, Peru, Surinam, Trinidad, Venezuela; Belize, Costa Rica (incl. Cocos Island), El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama; Antigua, Barbados, Cuba, Dominica, Dominican Republic, Guadeloupe, Haiti, Hispaniola, Jamaica, Martinique, Montserrat, Nevis, Puerto Rico (incl. Mona Is., Vieques Is.), St. Barthelemy, St. Croix, St. Kitts, St. Lucia, St. Thomas, St. Vincent; NC, NE, SE, SW USA (Alonso-Zarazaga and Lyal 1999).

Distribution in Colombia. Representatives of the tribe have been recorded from 21 Departments in the Country: Amazonas, Antioquia, Bolívar, Boyacá, Caldas, Casanare, Cauca, Chocó, Cundinamarca, Magdalena, Meta, Nariño, Norte de Santander, Putumayo, Quindío, Risaralda, Santander, Tolima, Valle del Cauca, Vaupés and Vichada (Suppl. material 2).

Bibliographic resources. The *Biologia Centrali-Americana* (Sharp and Champion 1911) may be a good place to start understanding the diversity of Eustylini (as ‘*Exophthalmina*’, p. 247 and ‘*Playomina*’, p. 282). There is a key to identify species of *Brachyomus* in Faust (1893, 16, in German). Marshall (1922b) discusses several classification issues with *Compsus*, *Exophthalmus* Schönherr, 1823 (Schönherr 1823, col. 1140) and allies, describes new genera and species and presents a key to identify species of *Exorides* (p. 203). Hustache (1938a, in French) revised part of South American *Compsus*; O’Brien and Peña (2012) translated to English and modified Hustache’s key and provided identification and records for two species of *Compsus* recorded as *Citrus* pests in Colombia. Franz (2010a) re-described several type species in Eustylini and presented a morphology-based phylogeny of the so called “*Exophthalmus* genus complex”, which involves several eustyline genera, especially Central American and Caribbean groups. Girón and Chamorro (2020) defined the “*Compsus* genus complex” and discussed affinities amongst species in this group. Some eustyline genera were included in phylogenetic analyses emphasising South American taxa by Marvaldi et al. (2018).

Remarks. The circumscription of Eustylini has been problematic since Lacordaire (1863), in French (some as ‘*Cyphides*’ p. 107, some as ‘*Geonémides*’ p. 125 and some as ‘*Eustylides*’ p. 205). Different genera, currently placed in Eustylini, have also been grouped together with members of Phyllobiini and Naupactini through time; most genera were treated as *incertae sedis* in the *Coleopterorum Catalogus* (Lona 1938: 508–525, 530–532). Kuschel (in Wibmer and O’Brien 1986a) brought together the majority of the genera that now compose Eustylini into a single tribe; the tribe is treated as ‘*Eustylina*’ by Morrone (1999). Some additions to the tribe have been made with transfers from other tribes, justified by a morphology-based

phylogeny presented by Franz (2012). Marvaldi et al. (2018) made the most recent addition by placing *Galapagonotus* Anderson & Lanteri, 2000 (Anderson and Lanteri 2000, 3) in Eustylini; they indicate that *Coconotus* Anderson & Lanteri, 2000 (Anderson and Lanteri 2000, 6) might also be an eustyline, but this remains to be confirmed, as it also shares features with the geonemine genus *Lachnopus* Schönherr, 1840 (Schönherr 1840b, 380), particularly with species in the luctuosus species-group (see Girón et al. 2018). Recent phylogenetic analyses (Franz 2012; Zhang et al. 2017; Marvaldi et al. 2018) indicate close relationships of Eustylini with taxa currently placed in Geonemini Gistel, 1856 (Gistel 1856, 373), which is primarily Caribbean and Central American.

The generic limits between some eustyline genera are not clearly defined; this is particularly true for *Compsus*, *Oxyderces* Schönherr, 1823 (Schönherr 1823, col. 1140), *Exorides* and *Xestogaster* Marshall, 1922 (Marshall 1922b, 221), all well represented in Colombia. This highlights the pressing need for taxonomic and phylogenetic revisions in the tribe, including (ideally recent) Colombian material that is already deposited in national collections. This particular group is part of the “*Compsus* genus complex” (Girón and Chamorro 2020); the authors compare several species of *Compsus*, *Oxyderces* and *Exophthalmus* which, by their colouration, may be confused with the golden-headed weevil, *Compsus auricephalus* (Say, 1824) (Say 1824), which is the only species of *Compsus* to occur in North America.

Members of Eustylini are amongst the most commonly found entimines in Colombian biological collections; *Compsus* is the most frequently observed and collected genus, followed by *Exorides* and perhaps *Eustylus*. Species of *Exophthalmus* are rare and seem to be restricted to the Pacific Region. Species of *Compsus* have been recorded as *Citrus* pests (Cano et al. 2002a, b; Gallego et al. 2012, O'Brien and Peña 2012) in Colombia and sugar cane in Venezuela (Kuschel 1955a). The Colombian endemic *Oxyderces viridipes* (Boheman in Schönherr 1840b, 179) has been intercepted at ports in the USA on *Hydrangea* flowers (Hydrangeaceae; see Girón and Chamorro 2020). In nature, eustylinines commonly rest with their prolegs extended towards the front of the body.

Tribe Lordopini Schönherr, 1823

Fig. 8

Recognition. Medium-sized weevils (approx. 10–20 mm); scale coverage variable in presence, density, pattern and colouration; iridescent scales and setae may be present; surface usually even (not coarsely sculptured); head (including rostrum, Fig. 8B, C) subrectangular to conical, usually longer than wide; eyes usually large and elongated in lateral view, weakly to moderately projecting from surface of head; frons nearly as wide as or wider than interantennal distance, often bearing median fovea; rostrum nearly parallel-sided, broadened apically (Fig. 8B) or with lateral margins apically converging (Fig. 8C); dorsal surface of rostrum either flat




Figure 8. Distribution and some head variants of Lordopini: **A** distribution map of records for *Hypsonotus* and undetermined lordopines in Colombia **B**, **C** head in dorsal view: **B** *Hypsonotus* **C** *Lordops* **D** head and anterior section of prothorax in lateral view of *Hypsonotus* showing well-developed postocular lobe with setae forming a fringe.

or with one or three longitudinal carinae; antennal scrobe generally only visible at apical region in dorsal view; nasal plate (see Fig. 1B, C) usually well developed, flat or depressed regarding surface of rostrum; anterior margin of prothorax in lateral view strongly sinuate, forming conspicuous postocular lobe (Fig. 8D); postocular setae present, forming a fringe instead of a tuft (Fig. 8D); elytral shoulders absent to well-developed (see Fig. 1N–P); tubercles may be present on elytra; legs relatively slender.

Amongst entimines with well-developed postocular lobe, lordopines are recognisable because of their elongated rostrum (Fig. 8B, D) or their conical heads with apically-converging lateral margins of the rostrum (Fig. 8C). They may be

confused with mid-sized Entimini, but the bulky shape of the body of most Entimini and the overall shape of the head differentiate them (compare Fig. 5B, C with Fig. 8B, C).

Diversity. There are 44 genera with 291 species of Lordopini described (Wibmer and O'Brien 1986a). Eleven genera, represented by 33 species, have been recorded from Colombia, four and 25, respectively, being endemic.

General distribution. Lordopini is a strictly Neotropical tribe, ranging from Mexico to Paraguay, including some of the Lesser Antilles. Lordopies have been recorded from the following countries: Argentina, Bolivia, Brazil, Colombia, Ecuador, French Guiana, Paraguay, Peru, Venezuela; Costa Rica, Belize, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama; Grenada, St. Vincent (Alonso-Zarazaga and Lyal 1999). The higher diversity of this tribe is massively concentrated in Brazil.

Distribution in Colombia. Representatives of Lordopini have been recorded from Bolívar, Magdalena, Nariño, Quindío, Santander, Valle del Cauca and Vaupés (Suppl. material 2).

Bibliographic resources. A general diagnosis for Lordopini can be found in Lacordaire (1863, 260, as 'Hypsonotides', in French). Only two genera appear in the Biologia Centrali-Americana (Sharp and Champion 1911, 300, under 'Entimina'). Kessel (1932, 1935, 1937, in German) revised several genera and provided keys to genera and species, focusing on Brazilian fauna, but these volumes are not yet available online and are rather difficult to find in libraries.

Remarks. Lordopini is one of the least-studied entimine tribes in the region and needs systematic revision: 18 genera are Brazilian endemics; more than one third of the species belong to *Hypsonotus* Germar, 1824 (Germar 1824, 367, 115 spp.) and one fifth belongs to the genus *Lordops* Schönherr, 1823 (Schönherr 1823, col. 1142, 59 spp.). From the remaining genera, 17 are monotypic and 11 are represented by only two species. All of these facts might indicate problems with current generic definitions. Furthermore, there are no phylogenetic analyses, including lordopines, to date. There are records of an undetermined species of *Hypsonotus* attacking peanuts in Brazil (Araujo et al. 1977); other than that, the natural history of the Lordopini remains largely unknown.

Tribe Naupactini Gistel, 1856

Fig. 9

Recognition. Small to medium weevils (approx. 3.5–35 mm); scale coverage highly variable in presence, density and colouration; iridescent scales, erect setae or waxy secretions are frequently present; surface usually even; head (including rostrum; Fig. 9C–F) subrectangular to trapezoid, nearly as long or shorter than wide; eyes rounded to oval in lateral view, small to large, laterally positioned and moderately to strongly projected from surface of head; frons usually as wide as or wider than interantennal distance, often bearing median longitudinal furrow extending through

rostrum; rostrum with lateral margins usually apically converging or nearly parallel-sided, sometimes slightly broadened apically (Fig. 9C–F); dorsal surface of rostrum usually flat (excepting median furrow; broadly anteriorly depressed in *Platyomus* Sahlberg, 1823 (Sahlberg 1823, 29) and relatives, for example, Fig. 9F); antennal scrobe generally not broadly visible in dorsal view; nasal plate (see Fig. 1B, C) of variable development, usually flat to depressed regarding surface of rostrum; anterior margin of prothorax in lateral view straight, never forming postocular lobe; postocular setae absent; elytral shoulders absent to well-developed (see Fig. 1N–P); tubercles may be present on elytra; profemora sometimes enlarged (see Fig. 1J); apical region of metatibia rectangular, fringed by spines along ventral and posterior margins (Fig. 1M).

Naupactines are relatively easy to recognise amongst South American entimines by the overall shape of their head, the usual presence of a median longitudinal furrow along the head and rostrum and the characteristic apex of their metatibia. They may be confused with similarly-coloured Eustylini, but the overall shape of the head (compare Fig. 7C–F with Fig. 9C–F) and the apex of the metatibia differentiate them (compare Fig. 7C–F with Fig. 9C–F). Small naupactines may be confused with anypotactines, from which they can be differentiated by the metatibial apex. From Tanymecini, naupactines can be distinguished by the lack of postocular setae in Naupactini and the shape of the metatibial apex.

Diversity. Naupactini contains 67 genera with over 500 species described (Lanteri and del Río 2016a). Fourteen genera, represented by 36 species have been recorded from Colombia, two genera and 22 species being endemic.

General distribution. In the New World, Naupactini ranges from the USA to Argentina and Chile, including some of the Caribbean islands. Naupactines have been recorded from the following countries: Argentina, Bolivia, Brazil, Chile (including Isla de Pascua, Juan Fernández Is.), Colombia, Ecuador (including Galapagos Is.), French Guiana, Guyana, Paraguay, Peru, Uruguay, Surinam, Trinidad, Venezuela; Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama; USA (Alonso-Zarazaga and Lyal 1999). Its highest diversity is concentrated in Argentina, Bolivia, Brazil, Paraguay and Uruguay, which coincides with areas where they have been extensively studied.

Distribution in Colombia. Representatives of the tribe have been recorded from 18 Departments: Antioquia, Bolívar, Boyacá, Caldas, Casanare, Cauca, Cesar, Chocó, Cundinamarca, Guaviare, Magdalena, Meta, Nariño, Quindío, Risaralda, Valle del Cauca, Vaupés and Vichada (Suppl. material 2).

Bibliographic resources. A recent summary of the status of knowledge in Naupactini can be found in Lanteri and del Río (2016a). Several genera of Naupactini have been revised or described since the 1980s (Lanteri 1980, 1982, 1984, 1985, 1986, 1989, 1990a, b, c, d, 1992, 1995; Lanteri and Loíacono 1990; Lanteri and O'Brien 1990; Bordón 1991, 1997; Lanteri and Morrone 1991, 1995; Lanteri and Díaz 1994; Lanteri and Marvaldi 1995; Lanteri and del Río 2003, 2004, 2005, 2006a, a, 2016b; del Río et al. 2006, 2012; del Río and Lanteri 2007a, b, 2011a, b,




Figure 9. Distribution and some head variants of Naupactini: **A, B** distribution maps of Naupactini in Colombia: **A** records for *Amphideritus*, *Lanterius*, *Mimographus* and *Naupactus* **B** records for *Litostylus*, *Platyomus* and undetermined naupactines **C–F** head in dorsal view: **C** *Mimographus* **D** undetermined Naupactini **E** *Litostylus* **F** “*Platyomus*” (see discussion under Naupactini) with characteristic broad and flattened antennal scape.

2012, 2013, 2019, 2020; Lanteri et al. 2017). There have also been recent efforts to elucidate the phylogenetic relationships amongst naupactine genera (Lanteri and del Río 2016a; del Río et al. 2018) and with other entimines (Marvaldi et al. 2018). Keys to genera can be found in Sharp and Champion (1911, 221) and van Emden (van Emden 1944b); a key to Andean genera is available in del Río and Lanteri (2011a).

Remarks. Naupactini is the largest tribe of New World entimines and one of the best known, mainly because of the revisionary and phylogenetic efforts of Dr. Analía Lanteri, Dr. Adriana Marvaldi, Dr. Guadalupe del Río and collaborators, based in Argentina. Different naupactine clades are specific to certain biogeographic regions in South America (see Lanteri 1992; del Río and Lanteri 2011a), with flightlessness

and parthenogenesis being frequent in certain systems (Lanteri and Normark 1995; Guzmán et al. 2012; Lanteri et al. 2013; Lanteri and del Río 2016a). A detailed study of the female genitalia and oviposition habits in Naupactini, in a phylogenetic context, can be found in Lanteri and del Río (2008).

Most revisionary efforts to date have focused on fauna from Argentina, Brazil and surrounding countries. In northern South America, naupactines from Venezuela were studied by Bordón [1991, genus *Macrostylus* Boheman, 1840 (Schönherr 1840a, 921), 26 spp. described as new; 1997, genus *Naupactus* Dejean, 1821 (Dejean 1821, 94), 24 spp. described as new]; none of these species has been recorded from Colombia to date. Other than that, the naupactine fauna from northern South America, Central America and the Caribbean is still in need of revision.

Several naupactine species are economically important (e.g. Lanteri et al. 2002, 2013); *Leschenius vulcanorum* (Kirsch, 1889) (Kirsch 1889, 17) has been recorded in Colombia as the ‘Potato shooter’ or ‘tiroteador de la papa’ (Canchala 1992; Peña 2001; Cortázar Gómez et al. 2012).

Members of Naupactini are very common in Colombian entomological collections, especially the genera *Lanterius* Alonso-Zarazaga & Lyal, 1999 (Alonso-Zarazaga and Lyal 1999, 164), *Mimographus* Schönherr, 1847 (Schönherr 1847, 34) and allies. *Platyomus* was recorded for the first time for Colombia by Girón (2007a; for Risaralda and Valle del Cauca) and then by Cardona-Duque et al. (2018; for Antioquia); these records might be erroneously identified and, instead, the specimens cited could be representatives of *Chamaelops* Kirsch, 1868 (Kirsch 1868, 235), which is fairly closely related to *Platyomus* (Lanteri and del Río 2016a); *Chamaelops* has previously been recorded for Colombia, considered endemic and includes two described species. *Galapaganus* Lanteri, 1992 (Lanteri 1992, 230) and *Pantomorus* Schönherr, 1840 (Schönherr 1840a, 942) were also recorded by Girón (2007a; both from Valle del Cauca) and are presented here (see Suppl. material 2; including records from Bolívar for *Galapaganus* and from Casanare and Meta for *Pantomorus*), but these records need to be confirmed.

Lanterius was erected by Alonso-Zarazaga and Lyal (1999, 164) to group species that were previously placed in a misidentified concept of *Mimographus* Schönherr, 1847 (Schönherr 1847, 34; see del Río and Lanteri 2018, 331). Even though del Río and Lanteri (2018) indicate that each of the *Mimographus* species, listed by Wibmer and O’Brien (1986a, 56), needs to be studied in detail in order to determine their appropriate placement, the Colombian species are here listed as new combinations under *Lanterius* in an attempt to prevent confusion, as there are species of true *Mimographus* recorded from Colombia as well. These new combinations are: *Lanterius ardosiacus* (Kirsch, 1868: 229), comb. nov., *Lanterius cinereoguttatus* (Champion, 1911: 228), comb. nov., *Lanterius hirtus* (Voss, 1953: 60), comb. nov., *Lanterius laesicollis* (Schönherr, 1847: 35), comb. nov., *Lanterius micans* (Kirsch, 1868: 229), comb. nov., *Lanterius rotundicollis* (Kirsch, 1889: 15), comb. nov., *Lanterius versicolor* (Kirsch, 1889: 15), comb. nov., and *Lanterius vittatus* (Kirsch, 1889: 14), comb. nov.

Tribe Premnotrypini Kuschel, 1956

Fig. 10D

Recognition. Small weevils (approx. 4–15 mm); scale coverage usually dense, uniform, and brown, often with patches of erect scales or setae; surface from even to undulate or tuberculate; head (including rostrum, Fig. 10D) usually conical and longer than wide; eyes laterally positioned and weakly to moderately projected from surface of head; frons wider than interantennal distance; rostrum elongate, usually with lateral margins apically converging, sometimes slightly broadened apically; dorsal surface of rostrum flat to undulate; mandibular scar reduced to absent; antennal scrobe generally not broadly visible in dorsal view; nasal plate (see Fig. 1B, C) of variable development, usually flat to elevated regarding surface of rostrum; anterior margin of prothorax in lateral view sinuate, forming well-developed and large postocular lobe (see Fig. 1F); postocular setae forming fringe; elytral shoulders usually well-developed (see Fig. 1P); tubercles may be present on elytra.

Premnotrypines may not be easily recognisable as entimines, as their rostrum is relatively slender and usually directed towards the body when mounted; the reduction or absence of mandibular scars would make them pass as other miscellaneous Curculionidae. Amongst entimines, they can be recognised by their small size and well-developed postocular lobe, fringed by postocular setae. By their size, premnotrypines can be confused with anypotactines or small tanymecines, but can be recognised by the presence of a well-developed postocular lobe, lacking in both Anypotactini and Tanymecini.

Diversity. There are only three genera in Premnotrypini with 28 species (Kuschel 1956; Wibmer and O'Brien 1986a, 88). There is only one species recorded from Colombia.

General distribution. Premnotrypini is an exclusively Andean group, distributed from Colombia to Bolivia and northern Chile. Their diversity is concentrated in Bolivia and Peru, at elevations over 3000 m a.s.l. No specimens of this tribe were identified in the collections revised for this work.

Bibliographic resources. The tribal diagnosis, key to genera, descriptions of two of the genera and keys to all species can be found in Kuschel (1956, in Spanish).

Remarks. The genus *Premnotrypes* is associated with potato cultivars and contains economically-important species commonly known as the “Andean potato weevils” (see Alcázar and Cisneros 1998). There are many studies in the natural history and control of pest species (e.g. Alcalá and Alcázar 1976; Calvache 1987; Dueñas 1989), but no phylogenetic studies in Premnotrypini have been published to date.

Tribe Tanymecini Lacordaire, 1863

Fig. 10A–C

Recognition. Small to large weevils (approx. 4–20 mm); scale coverage variable in presence, density and colouration, often dense and uniform, with colouration

patterns formed by scales on dorsal surface; surface usually even, sometimes granulate; head (including rostrum; Fig. 10B, C) usually subrectangular and nearly as long as wide; eyes laterally positioned and weakly to strongly projected from surface of head; frons usually as wide as interantennal distance, bearing median fovea or small (narrow and short) longitudinal furrow that may extend along rostrum; rostrum subquadrate, usually with lateral margins parallel, sometimes slightly constricted basad of antennal insertion; dorsal surface of rostrum flat to slightly depressed; antennal scrobe only visible along anterior section in dorsal view; epistoma usually well-developed; nasal plate (see Fig. 1B, C) of variable development, flat to depressed regarding surface of rostrum; anterior margin of prothorax in lateral view straight (Fig. 1D), never forming well-developed postocular lobe; postocular setae usually present, clearly visible, grouped as a tuft, usually positioned near ventral margin of prothorax (Fig. 1D); elytral shoulders absent to well-developed (see Fig. 1N–P); tubercles may be present on elytra and elytral apices may be projected; profemora frequently enlarged (see Fig. 1J).

Amongst Neotropical entimines, members of Tanymecini are easily recognisable by the presence of postocular setae, although not all tanymecines have this feature and not all the species with postocular setae are tanymecines. The postocular setae in Tanymecini are usually clustered, forming a tuft as opposed to distributed along the margin forming a fringe (compare Fig. 1D with Fig. 1F). Some tanymecines can be potentially confused with Naupactini because of their flat frons, median furrow and laterally-positioned eyes, but the presence of postocular setae in Tanymecini sets them apart easily; they can also be differentiated by the particular shape of the apex of the metatibia in Naupactini, which is rectangular and fringed by spines along the ventral and posterior margins (Fig. 1M; compare with rounded margins in tanymecines, Fig. 1L).

Diversity. There are about 100 genera of Tanymecini distributed worldwide. Eighteen genera are represented throughout the Americas by 325 species (O'Brien and Wibmer 1982, 45; Wibmer and O'Brien 1986a, 67; Alonso-Zarazaga and Lyal 1999, 178; Morrone 1999, 143; Cortés-Hernández and Anderson 2019; Cortés-Hernández and Morrone 2020). There are 58 species of Tanymecini recorded from Colombia, 29 of them being considered endemic.

General distribution. Tanymecines are distributed throughout the Americas and the Caribbean islands, including the following countries: Argentina, Bolivia, Brazil, Chile, Colombia, Ecuador, Peru, Paraguay, Uruguay, Venezuela; Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama; Bahamas, Cuba, Dominica, Dominican Republic, Grand Cayman, Grenada, Guadeloupe, Haiti, Jamaica, Saint Vincent; C, E Canada, USA (Alonso-Zarazaga and Lyal 1999). Their highest diversity is concentrated in Central America and northern South America.

Distribution in Colombia. Tanymecines have been found in collections from Antioquia, Bolívar, Boyacá, Caldas, Cauca, Cundinamarca, Magdalena, Meta, Nariño, Norte de Santander, Quindío, Risaralda and Valle del Cauca (Suppl. material 2).




Figure 10. Distribution and some head variants of Tanymecini and Premnotrypini: **A** distribution map of records for *Hadromeropsis*, *Macropterus*, *Pandeleteius* and undetermined tanymecines in Colombia **B**, **C** head in dorsal view of Tanymecini: **B** *Pandeleteius* **C** *Hadromeropsis* **D** head in dorsal view of *Premnotrypes* (Premnotrypini).

Bibliographic resources. Most genera of Tanymecini represented in the Americas were revised by Anne Howden (1959, 1961, 1963, 1966, 1969, 1970, 1974, 1976, 1982, 1986, 1993, 1996, 1998, 2001, 2004, 2008, 2011; Girón and Howden 2019). A key to genera recorded from Mexico was published by Cortés-Hernández and Morrone (2019), which includes most genera represented in Colombia. Howden's revision of the species of *Pandeleteius* of Colombia and Venezuela (Howden 1976) considerably increased the number of species known from Colombia and includes a key to species.




Figure 11. Distribution of undetermined entimines in Colombia.

Remarks. Tanymecini is one of the best-known tribes of entimines of the Americas, thanks to the works of Anne Howden. Tanymecines, especially *Pandeteleius*, are quite common in Colombia. In nature, tanymecines are commonly seen with their prolegs resting perpendicular to the body axis, a type of behaviour shared with some naupactines.

List of valid species of Entiminae recorded from Colombia

Genera and species in bold letters indicate endemic taxa. Fossil taxa are indicated by (+). Localities in bold letters indicate type locality. Localities marked with an asterisk are records taken from ASUCOB. A double asterisk means that identifications need to be corroborated. A question mark (?) in the distribution indicates that the locality record is doubtful and needs confirmation. Type localities from countries other than Colombia are omitted. Taxon names include author, year and page number of the original description (e.g. *Anypotactus* Schönherr 1840b, 299); for some of Schönherr's taxa, the page number corresponds to a column number in the publication and is indicated by 'col.' (e.g. Entiminae Schönherr 1823, col. 1138). See other locality remarks in the methods section.

Tribe Anypotactini Champion, 1911: 215

Genus *Anypotactus* Schönherr, 1840: 299

Distribution: Bolivia, Colombia, Peru, Venezuela; Costa Rica, El Salvador, Guatemala, Honduras, Nicaragua, Panama.

References: Schönherr 1840b, 299; Sharp and Champion 1911, 215, 174; Kuschel 1955b, 275, O'Brien and Wibmer 1982, 29; Wibmer and O'Brien 1986a, 44; Alonso-Zarazaga and Lyal 1999, 145; Morrone 1999, 135.

Anypotactus exilis Boheman, 1840: 300

Distribution: **Colombia**, Venezuela; Costa Rica, Guatemala, Honduras, Nicaragua, Panama.

References: Boheman in Schönherr 1840b, 300; Sharp and Champion 1911, 216, O'Brien and Wibmer 1982, 29; Wibmer and O'Brien 1986a, 44; Morrone 1999, 135.

Anypotactus morosus (Boheman, 1840: 449)

Distribution: **Colombia**.

References: As *Polydrosus morosus* by Boheman in Schönherr 1840b, 449; Kuschel 1955b, 277 (as *Anisactus morosus*), Kuschel in Wibmer and O'Brien 1986a, 44; Morrone 1999, 135.

Genus *Bothinodontes* Kirsch, 1868: 241

Distribution: Colombia (Caldas, Tolima); Mexico (?).

References: Kirsch 1868, 241; Gemmiger and von Harold 1871, 2242; Sharp and Champion 1911, 331, O'Brien and Wibmer 1982, 29; Wibmer and O'Brien 1986a, 45; Alonso-Zarazaga and Lyal 1999, 145; Morrone 1999, 135.

Bothinodontes gibbipennis (Champion, 1911: 332)

Distribution: Colombia; **Mexico** (?).

References: As *Bothynodontes* by Champion in Sharp and Champion 1911, 332, O'Brien and Wibmer 1982, 29; Wibmer and O'Brien 1986a, 45; Morrone 1999, 135.

***Bothinodontes squalidus* Kirsch, 1868: 243**

Distribution: Colombia ('**Bogotá**').

References: Kirsch 1868, 243; Wibmer and O'Brien 1986a, 45; Morrone 1999, 135.

Genus *Cylloproctus* Faust, 1892: 22

Distribution: Brazil, Colombia, Venezuela.

References: Faust 1892, 22; Wibmer and O'Brien 1986a, 45; Alonso-Zarazaga and Lyal 1999, 145; Morrone 1999, 135.

***Cylloproctus modestus* Faust, 1892: 23**

Distribution: **Colombia** [Columbia].

References: Faust 1892, 23 (in footnote), Wibmer and O'Brien 1986a, 45; Morrone 1999, 135.

***Cylloproctus pantomoroides* Voss, 1936: 109**

Distribution: Colombia (Cundinamarca: **Fusagasugá**).

References: Voss 1936, 109; Wibmer and O'Brien 1986a, 45; Morrone 1999, 135.

Genus *Hypsometopus* Kirsch, 1868: 222

Distribution: Colombia.

References: Kirsch 1868, 222; Wibmer and O'Brien 1986a, 45; Alonso-Zarazaga and Lyal 1999, 146; Morrone 1999, 136.

***Hypsometopus inquinatus* Kirsch, 1868: 223**

Distribution: Colombia ('**Bogotá**').

References: Kirsch 1868, 223; Wibmer and O'Brien 1986a, 45; Morrone 1999, 136.

Genus *Phanasora* Pascoe, 1881: 38

Distribution: Colombia, Ecuador.

References: Pascoe 1881, 38; Wibmer and O'Brien 1986a, 45; Alonso-Zarazaga and Lyal 1999, 146; Morrone 1999, 137.

***Phanasora plumbea* Pascoe, 1881: 39**

Distribution: Colombia ('**Bogotá**').

References: Pascoe 1881, 39; Wibmer and O'Brien 1986a, 45; Morrone 1999, 137.

Genus *Polydacrys* Schönherr, 1834: 130

Distribution: Bolivia, Colombia, Cuba, Dominica, Granada, Guadeloupe, Puerto Rico, Saint Vincent; Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama; SW USA (Texas).

References: Schönherr 1834, 130; Sharp and Champion 1911, 217, O'Brien and Wibmer 1982, 29; Wibmer and O'Brien 1986a, 44; Alonso-Zarazaga and Lyal 1999, 146; Morrone 1999, 137.

***Polydacrys seriegranosus* Champion, 1911: 218**

Distribution: Bolivia, Colombia; Panama.

References: Sharp and Champion 1911, 218, O'Brien and Wibmer 1982, 29; Wibmer and O'Brien 1986a, 44; Morrone 1999, 137.

Genus *Prepodellus* Kirsch, 1868: 239

Distribution: Colombia; Costa Rica, Mexico, Nicaragua, Panama.

References: Kirsch 1868, 239, O'Brien and Wibmer 1982, 29; Wibmer and O'Brien 1986a, 44; Alonso-Zarazaga and Lyal 1999, 146; Morrone 1999, 137.

***Prepodellus nigriclavis* Kirsch, 1868: 240**

Distribution: Colombia ('**Bogotá**', Cundinamarca (La Vega*)).

References: Kirsch 1868, 240; Wibmer and O'Brien 1986a, 44; Morrone 1999, 137.

***Prepodellus ruficornis* Kirsch, 1868: 240**

Distribution: Colombia ('**Bogotá**', Cundinamarca (Mosquera*)).

References: Kirsch 1868, 240; Wibmer and O'Brien 1986a, 44; Morrone 1999, 137.

Tribe Entimini Schönherr, 1823: col. 1138

Genus *Cydianerus* Schönherr, 1840: 737

Distribution: Argentina, Bolivia, Brazil, Colombia, Paraguay, Venezuela; Costa Rica, El Salvador, Honduras, Mexico, Panama.

References: Schönherr 1840a, 737; Sharp and Champion 1911, 300 as Entimina, O'Brien and Wibmer 1982, 67; Wibmer and O'Brien 1986a, 101; Alonso-Zarazaga and Lyal 1999, 144; Morrone 1999, 110; Poinar et al. 2017 (key to species).

***Cydianerus eukrinus* Poinar, Bukejs & Legalov, 2017: 88 (+)**

Distribution: Colombia (**Santander**, copal resin).

References: Poinar et al. 2017, 88.

Cydianerus pascoei (Bovie, 1908a: 4)

Distribution: **Colombia**; Panama.

References: Bovie 1908a, 4 as *Polydius*, in footnote, O'Brien and Wibmer 1982, 67; Wibmer and O'Brien 1986a, 101; Morrone 1999, 111.

Genus *Entimus* Germar, 1817: 341

Distribution: Argentina, Bolivia, Brazil, Colombia, Ecuador, French, Guiana, Guyana, Paraguay, Peru, Uruguay; Costa Rica, Nicaragua, Panama.

References: Germar 1817, 341; Sharp and Champion 1911, 301, O'Brien and Wibmer 1982, 67; Wibmer and O'Brien 1986a, 99; Alonso-Zarazaga and Lyal 1999, 144; Morrone 1999, 111, 2002; Morrone et al. 2019.

Entimus arrogans Pascoe 1872: 448

Distribution: Colombia; Costa Rica, Nicaragua, **Panama**.

References: Pascoe 1872, 448; Sharp and Champion 1911, 301, O'Brien and Wibmer 1982, 67; Wibmer and O'Brien 1986a, 100; Morrone 1999, 111, 2002, 505.

Entimus granulatus (Linnaeus, 1758: 386)

Distribution: Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Peru; Panama.

References: Linnaeus 1758, 386 (as *Curculio granulatus*), O'Brien and Wibmer 1982, 67; Wibmer and O'Brien 1986a, 100; Morrone 1999, 111, 2002, 507.

Genus *Rhigus* Schönherr, 1823: col. 1138

Distribution: Argentina, Brazil, Colombia, French Guiana, Paraguay, Peru.

References: Schönherr 1823, col. 1138; Wibmer and O'Brien 1986a, 101; Alonso-Zarazaga and Lyal 1999, 145; Morrone 1999, 121; Gaiger 2001.

Rhigus speciosus (Linnaeus, 1758: 385)

Distribution: Brazil, Colombia (Amazonas, Tolima), French Guiana, Peru.

References: Linnaeus 1758, 385 (as *Curculio speciosus*), Wibmer and O'Brien 1986a, 102; Morrone 1999, 121; Gaiger 2001, 63; Girón and Cardona-Duque 2018, 181 (first record for Colombia), Gillett and Barr 2020 (first record for Peru).

Tribe Eudiagogini LeConte, 1874: 454

Genus *Colecerus* Schönherr, 1840: 927

Distribution: Colombia; Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua; SW USA.

References: Schönherr 1840a, 927; Sharp and Champion 1911, 305, O'Brien and Wibmer 1982, 58 (under Promecopini), Wibmer and O'Brien 1986a, 81 (under Promecopini), Alonso-Zarazaga and Lyal 1999, 157; Morrone 1999, 110 (under Entimina).

Colecerus albidus (Chevrolat, 1881: XXXVIII)

Distribution: Colombia ('**Bogotá**').

References: Chevrolat 1881, XXXIII, Wibmer and O'Brien 1986a, 81; Morrone 1999, 110.

Genus *Promecops* Sahlberg, 1823: 30

Distribution: Argentina, Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Paraguay, Peru, Venezuela; Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama; Grenadines, Guadeloupe, St. Vincent.

References: Sahlberg 1823, 30, O'Brien and Wibmer 1982, 59 (under *Promecopini*), Wibmer and O'Brien 1986a, 82 (under *Promecopini*), Alonso-Zarazaga and Lyal 1999, 157; Morrone 1999, 119 (under *Entimina*).

***Promecops lepidoides* Voss, 1934: 75**

Distribution: Colombia (Meta: **Villavicencio**).

References: Voss 1934, 75 (in key), Wibmer and O'Brien 1986a, 83; Morrone 1999, 120.

***Promecops lepidus* Fåhraeus, 1840: 317**

Distribution: **Colombia**, [Orinoco] (Magdalena (Ciénaga)*).

References: Chevrolat 1881, XXXIII, Wibmer and O'Brien 1986a, 81; Morrone 1999, 110.

Promecops leucothyreus Fåhraeus, 1840: 328

Distribution: Bolivia, Colombia ('**Bogotá**', Antioquia (Puerto Berrío), Bolívar (Cartagena)*, Magdalena); Guatemala, Honduras, Mexico, Nicaragua.

References: Fåhraeus in Schönherr 1840b, 328; Voss 1934, 80 (in key, as *Promecops leucothyrea*), O'Brien and Wibmer 1982, 59; Wibmer and O'Brien 1986a, 83; Morrone 1999, 120.

***Promecops rhombicus* Fåhraeus, 1840: 326**

Distribution: **Colombia** [Columbia].

References: Fåhraeus in Schönherr 1840b, 326; Voss 1934, 97 (listed as *Promecops rhombica*), Wibmer and O'Brien 1986a, 83; Morrone 1999, 120.

Promecops rhombifer Fåhraeus, 1840: 325

Distribution: **Colombia**, Guyana, Venezuela.

References: Fåhraeus in Schönherr 1840b, 325; Voss 1934, 90 (in key, as *Promecops rhombifera*), Wibmer and O'Brien 1986a, 83; Morrone 1999, 120.

Tribe Eustylini Lacordaire, 1863: 205

Genus *Brachyomus* Lacordaire, 1863: 130

Distribution: Colombia, Ecuador, French Guiana, Peru, Trinidad, Venezuela; St. Vincent.

References: Lacordaire 1863, 130; Wibmer and O'Brien 1986a, 73, 1986b; Alonso-Zarazaga and Lyal 1999, 158; Morrone 1999, 123.

Brachyomus quadrinodosus (Boheman, 1842: 217)

Distribution: Colombia, Venezuela.

References: Boheman in Schönherr 1842, 217 (as *Geonemus*), Wibmer and O'Brien 1986a, 74; Morrone 1999, 123.

Brachyomus quadrituberculatus (Boheman, 1842: 216)

Distribution: **Colombia**.

References: Boheman in Schönherr 1842, 216 (as *Geonemus*), Wibmer and O'Brien 1986a, 74; Morrone 1999, 123.

Genus *Compsus* Schönherr, 1823: col. 1140

Distribution: Argentina, Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Peru, Surinam, Venezuela; Costa Rica, Guatemala, Mexico, Nicaragua, Panama; Guadeloupe, Jamaica; NC, SE, SW USA.

References: Schönherr 1823, col. 1140; Hustache 1938a, O'Brien and Wibmer 1982, 54 (under Phyllobiini), Wibmer and O'Brien 1986a, 74; Alonso-Zarazaga and Lyal 1999, 158; Morrone 1999, 123, O'Brien and Peña 2012.

Compsus adonis Marshall, 1922: 197

Distribution: Colombia (Antioquia: **Medellín**).

References: Marshall 1922b, 197; Hustache 1938a, 71 (in key, as *Compsus bellus*, but see discussion in Girón and Chamorro 2020, 38), Wibmer and O'Brien 1986a, 74; Morrone 1999, 123, O'Brien and Peña 2012.

***Compsus aeruginosus* (Boheman, 1840: 177)**

Distribution: Colombia (**Antioquia**).

References: Boheman in Schönherr 1840b, 177 (as *Platyomus aeruginosus*), Wibmer and O'Brien 1986a, 74; Morrone 1999, 123.

***Compsus affinis* Hustache, 1938: 77**

Distribution: Colombia (Meta: **Villavicencio**).

References: Hustache 1938a, 77; Wibmer and O'Brien 1986a, 74; Morrone 1999, 123, O'Brien and Peña 2012, 4 (in key).

***Compsus albosetosus* Hustache, 1938: 84**

Distribution: **Colombia** (Cundinamarca: Bogotá).

References: Hustache 1938a, 84; Wibmer and O'Brien 1986a, 74; Morrone 1999, 124, O'Brien and Peña 2012, 5 (in key).

***Compsus albus* Hustache, 1938: 76**

Distribution: Colombia (Cundinamarca: Fusagasugá; Tolima: **Ibagué**).

References: Hustache 1938a, 76; Wibmer and O'Brien 1986a, 74; Morrone 1999, 124, O'Brien and Peña 2012, 4 (in key), Girón and Chamorro 2020, 39.

***Compsus alternevittatus* Hustache, 1938: 83**

Distribution: Colombia (**Rio Chili**, Boyacá (Muzo), Pensilvania (Caldas (?), Cundinamarca (?))).

References: Hustache 1938a, 83; Wibmer and O'Brien 1986a, 74; Morrone 1999, 124, O'Brien and Peña 2012, 5 (in key).

***Compsus ater* Kirsch, 1889: 20**

Distribution: Colombia (Pasto, **Santa Lucía**).

References: Kirsch 1889, 20; Wibmer and O'Brien 1986a, 74; Morrone 1999, 124.

***Compsus attenuatus* Hustache, 1938: 84**

Distribution: Colombia (Rio Chili; Boyacá: Muzo; Quindío Tolima: **Ibagué**).

References: Hustache 1938a, 84; Wibmer and O'Brien 1986a, 74; Morrone 1999, 124, O'Brien and Peña 2012, 5 (in key).

***Compsus bicarinatus* Kirsch, 1889: 22**

Distribution: Colombia (**Popayan**: Silvia (?)).

References: Kirsch 1889, 22; Wibmer and O'Brien 1986a, 74; Morrone 1999, 124.

***Compsus bituberculatus* Kirsch, 1889: 19**

Distribution: Colombia (**Cauca** [**Popayan**]: **Puracé**).

References: Kirsch 1889, 19; Wibmer and O'Brien 1986a, 74; Morrone 1999, 124.

***Compsus bituberosus* Kirsch, 1868: 237**

Distribution: Colombia (**Bogotá**).

References: Kirsch 1868, 237; Wibmer and O'Brien 1986a, 74; Morrone 1999, 124.

***Compsus candidus* Hustache, 1938: 75**

Distribution: Colombia (**Bogotá**); Valle del Cauca: Cartago, Palmira).

References: Hustache 1938a, 75; Wibmer and O'Brien 1986a, 74; Morrone 1999, 124, O'Brien and Peña 2012, 4 (in key).

***Compsus canescens* (Boheman, 1840: 181)**

Distribution: **Colombia**; Napo and Amazon rivers (Brazil (?), Ecuador (?), Peru (?)).

References: Boheman in Schönherr 1840b, 181 as *Platyomus canescens*, Guérin-Méneville 1855, 592 (listed), Wibmer and O'Brien 1986a, 75; Morrone 1999, 124.

Compsus cyanitarsis Hustache, 1938: 93

Distribution: Bolivia, Colombia.

References: Hustache 1938a, 93; Wibmer and O'Brien 1986a, 75; Morrone 1999, 124, O'Brien and Peña 2012, 6 (in key).

Compsus cyphoides Hustache, 1938: 108

Distribution: Colombia (Meta: **Villavicencio**).

References: Hustache 1938a, 108; Wibmer and O'Brien 1986a, 75; Morrone 1999, 124, O'Brien and Peña 2012, 6 (in key).

Compsus delicatulus Hustache, 1938: 79

Distribution: Colombia (Valle del Cauca: **Cartago**).

References: Hustache 1938a, 79; Wibmer and O'Brien 1986a, 75; Morrone 1999, 124, O'Brien and Peña 2012, 5 (in key).

Compsus deliciosus Hustache, 1938: 80

Distribution: Colombia (Risaralda: **Pereira**).

References: Hustache 1938a, 80; Wibmer and O'Brien 1986a, 75; Morrone 1999, 124, O'Brien and Peña 2012, 5 (in key).

Compsus deplanatus Kirsch, 1868: 237

Distribution: Colombia ('**Bogotá**'), Venezuela.

References: Kirsch 1868, 237; Wibmer and O'Brien 1986a, 75; Morrone 1999, 124.

Compsus divisus Hustache, 1938: 82

Distribution: Colombia (Boyacá: **Muzo**).

References: Hustache 1938a, 82; Wibmer and O'Brien 1986a, 75; Morrone 1999, 124, O'Brien and Peña 2012, 5 (in key), Girón and Chamorro 2020, 40.

Compsus eustylodes Hustache, 1938: 96

Distribution: Colombia (Cundinamarca: **Fusagasugá**, Viotá; Meta: Villavicencio).

References: Hustache 1938a, 96; Wibmer and O'Brien 1986a, 75; Morrone 1999, 124, O'Brien and Peña 2012, 3 (in key).

***Compsus glaucus* (Boheman, 1840: 178)**

Distribution: Colombia (**Antioquia**).

References: Boheman in Schönherr 1840b, 178 as *Platyomus glaucus*, Wibmer and O'Brien 1986a, 75; Morrone 1999, 125.

***Compsus iris* Marshall, 1922: 198**

Distribution: Colombia (Antioquia: **Medellín**).

References: Marshall 1922b, 198; Wibmer and O'Brien 1986a, 75; Morrone 1999, 125.

***Compsus latifrons* Hustache, 1938: 90**

Distribution: Colombia (**Pereira**).

References: Hustache 1938a, 90; Wibmer and O'Brien 1986a, 76; Morrone 1999, 125, O'Brien and Peña 2012, 4 (in key).

***Compsus lebasii* (Boheman, 1840: 182)**

Distribution: Colombia (Bolívar: **Cartagena**).

References: Boheman in Schönherr 1840b, 182; Hustache 1938a, 81; Wibmer and O'Brien 1986a, 76; Morrone 1999, 125, O'Brien and Peña 2012, 3 (in key).

***Compsus lineatus* Hustache, 1938: 81**

Distribution: Colombia (**S. Antonio**).

References: Hustache 1938a, 81; Wibmer and O'Brien 1986a, 76; Morrone 1999, 125, O'Brien and Peña 2012, 3 (in key).

***Compsus obliquatus* Hustache, 1938: 77**

Distribution: Colombia (**Rio Chili**; Cundinamarca: Beltrán; Tolima: Espinal).

References: Hustache 1938a, 77; Wibmer and O'Brien 1986a, 76; Morrone 1999, 125, O'Brien and Peña 2012, 4 (in key; p. 7 redescription).

Compsus pertinax Hustache, 1938: 85

Distribution: Colombia (**Cauca**), Ecuador.

References: Hustache 1938a, 85; Wibmer and O'Brien 1986a, 76; Morrone 1999, 125, O'Brien and Peña 2012, 4 (in key).

***Compsus placidus* (Boheman, 1840: 180)**

Distribution: **Colombia** [Nova Granata].

References: Boheman in Schönherr 1840b, 180 as *Platyomus placidus*, Wibmer and O'Brien 1986a, 76; Morrone 1999, 125.

***Compsus popayanus* Kirsch, 1889: 21**

Distribution: Colombia (**Popayán**).

References: Kirsch 1889, 21; Wibmer and O'Brien 1986a, 76; Morrone 1999, 125.

Compsus pugionatus Marshall, 1922: 196

Distribution: Colombia, Venezuela; Central America.

References: Marshall 1922b, 196; Wibmer and O'Brien 1986a, 76; Morrone 1999, 125.

Compsus quadrisignatus (Boheman, 1840: 188)

Distribution: Colombia, French Guiana, Venezuela.

References: Boheman in Schönherr 1840b, 188 as *Platyomus quadrisignatus*, Wibmer and O'Brien 1986a, 76; Morrone 1999, 125.

Compsus roseomicans Hustache, 1938: 100

Distribution: Colombia (Boyacá: Muzo).

References: Hustache 1938a, 100; Wibmer and O'Brien 1986a, 76; Morrone 1999, 125, O'Brien and Peña 2012, 6 (in key).

Compsus scrutator Hustache, 1938: 98

Distribution: Brazil, Colombia (**Cauca**), Venezuela.

References: Hustache 1938a, 98; Wibmer and O'Brien 1986a, 76; Morrone 1999, 125, O'Brien and Peña 2012, 6 (in key).

***Compsus violaceus* Hustache, 1938: 89**

Distribution: Colombia (**Rio Chili**).

References: Hustache 1938a, 89; Wibmer and O'Brien 1986a, 77; Morrone 1999, 126, O'Brien and Peña 2012, 5 (in key), Girón and Chamorro 2020, 40.

Compsus viridissimus Hustache, 1938: 107

Distribution: Colombia (Boyacá: **Muzo**), Ecuador.

References: Hustache 1938a, 107; Wibmer and O'Brien 1986a, 77; Morrone 1999, 126, O'Brien and Peña 2012, 6 (in key).

Compsus viridivittatus (Guérin-Ménéville, 1855: 592)

Distribution: Colombia (Antioquia (Támesis), Caldas (Chinchiná), Cundinamarca (Beltrán), Quindío (Armenia, La Tebaida, Montenegro), Risaralda (Belén de Umbria, Pereira), Santander (Lebrija), Tolima (Armero Guayabal, 4 Km SE Ibagué), Valle del Cauca (Caicedonia, Sevilla); **Napo and Amazon rivers** (Brazil (?), Ecuador (?), Peru (?)).

References: Guérin-Ménéville 1855, 592 as *Platyomus viridivittatus*, Wibmer and O'Brien 1986a, 77; Morrone 1999, 126, O'Brien and Peña 2012, 4 (in key; p. 10 re-description).

***Compsus viridulus* Hustache, 1938: 95**

Distribution: Colombia (**Bogotá**).

References: Hustache 1938a, 95; Wibmer and O'Brien 1986a, 77; Morrone 1999, 126, O'Brien and Peña 2012, 4 (in key).

Compsus zebra Marshall, 1922: 199

Distribution: Colombia (Valle del Cauca: Buenaventura), **Ecuador**.

References: Marshall 1922b, 199; Wibmer and O'Brien 1986a, 77; Morrone 1999, 126.

***Compsus zebrinus* Voss, 1953: 57**

Distribution: Colombia [West Cordillera. Rio Aguacatal, St. Antonio, Monte Socorro].

References: Voss 1953, 57; Wibmer and O'Brien 1986a, 77; Morrone 1999, 126.

Genus *Eustylus* Schönherr, 1842: 40

Distribution: Bolivia, Brazil, Colombia, French Guiana, Guyana, Paraguay, Peru, Trinidad, Venezuela; Costa Rica, Guatemala, Mexico, Panama; Guadeloupe, St. Lucia.

References: Schönherr 1842, 40; Sharp and Champion 1911, 290; Marshall 1916 (most Colombian species), Wibmer and O'Brien 1986a, 73; Alonso-Zarazaga and Lyal 1999, 158; Morrone 1999, 127.

Eustylus bodkini Marshall, 1916: 456

Distribution: Colombia, **Guyana**, Venezuela.

References: Marshall 1916, 456; Wibmer and O'Brien 1986a, 73; Morrone 1999, 127.

***Eustylus ehippiatus* Marshall, 1916: 458**

Distribution: **Colombia**.

References: Marshall 1916, 458; Wibmer and O'Brien 1986a, 73; Morrone 1999, 127.

***Eustylus magdalenae* Marshall, 1926: 536**

Distribution: Colombia (Magdalena: El Banco [El Blanco], Magdalena Valley; Aracataca).

References: Marshall 1926, 536; Wibmer and O'Brien 1986a, 73; Morrone 1999, 127.

Eustylus puber (Olivier, 1807: 367)

Distribution: Colombia, **French Guiana**, Guyana, Trinidad, Venezuela.

References: Olivier 1807, 367 (as *Curculio puber*), Wibmer and O'Brien 1986a, 73; Morrone 1999, 127; Franz 2010a, 56 (re-description).

***Eustylus simplex* Marshall, 1916: 460**

Distribution: **Colombia**.

References: Marshall 1916, 460; Wibmer and O'Brien 1986a, 73; Morrone 1999, 127.

***Eustylus simulatus* Marshall, 1916: 459**

Distribution: **Colombia**.

References: Marshall 1916, 459; Wibmer and O'Brien 1986a, 73; Morrone 1999, 127.

***Eustylus sordidus* Marshall, 1916: 460**

Distribution: **Colombia**.

References: Marshall 1916, 460; Wibmer and O'Brien 1986a, 73; Morrone 1999, 127.

Genus *Exophthalmus* Schönherr, 1823: col. 1140

Distribution: Brazil, Colombia, Ecuador, French Guiana, Surinam; Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama; Cuba, Dominican Republic, Guadeloupe, Haiti, Jamaica, Martinique, Puerto Rico.

References: Schönherr 1823, col. 1140; Sharp and Champion 1911, 249, O'Brien and Wibmer 1982, 56; Wibmer and O'Brien 1986a, 79; Alonso-Zarazaga and Lyal 1999, 158; Morrone 1999, 127; Franz 2012 (part).

Exophthalmus annulonotatus (Waterhouse, 1879: 423)

Distribution: Colombia (Antioquia: **Medellín**), Ecuador.

References: Waterhouse 1879, 423 (as *Paepodes annulonotatus*), Wibmer and O'Brien 1986a, 79; Morrone 1999, 128.

***Exophthalmus consobrinus* (Marshall, 1922b: 190)**

Distribution: Colombia (Chocó: Andagoya, Río Condoto; Valle del Cauca (Buenaventura)).

References: Marshall 1922b, 190 (as *Exophthalmodes consobrinus*), Wibmer and O'Brien 1986a, 79; Morrone 1999, 128.

Exophthalmus crassicornis Kirsch, 1868: 238

Distribution: Colombia (**Bogotá**); Panama.

References: Kirsch 1868, 238, O'Brien and Wibmer 1982, 56; Wibmer and O'Brien 1986a, 79; Morrone 1999, 128.

Exophthalmus jekelianus (White, 1858: 357)

Distribution: Colombia (**Chocó**); Costa Rica, Nicaragua, Panama.

References: White in Jekel 1858, 357 (as *Praepodes jekelianus*), O'Brien and Wibmer 1982, 57; Wibmer and O'Brien 1986a, 79; Morrone 1999, 128.

Exophthalmus sulcicrus Champion, 1911: 268

Distribution: Colombia (Chocó); Costa Rica, Guatemala, Nicaragua, Panama.

References: Sharp and Champion 1911, 268, O'Brien and Wibmer 1982, 58; Wibmer and O'Brien 1986a, 79; Morrone 1999, 129.

Genus *Exorides* Pascoe, 1881: 43

Distribution: Colombia, Ecuador, Peru, Venezuela.

References: Pascoe 1881, 43; Marshall 1922b, 202; Wibmer and O'Brien 1986a, 80; Alonso-Zarazaga and Lyal 1999, 158; Morrone 1999, 130.

Exorides bifurcatus Marshall, 1926: 537

Distribution: Colombia (Magdalena: **Sierra de San Lorenzo**).

References: Marshall 1926, 537; Wibmer and O'Brien 1986a, 80; Morrone 1999, 130.

Exorides caudatus Marshall, 1922: 217

Distribution: Colombia (**Bogotá**).

References: Marshall 1922b, 217; Wibmer and O'Brien 1986a, 80; Morrone 1999, 130.

Exorides cylindricus Marshall, 1922: 213

Distribution: Colombia (**Magdalena, San Lorenzo Mt.**).

References: Marshall 1922b, 213; Wibmer and O'Brien 1986a, 80; Morrone 1999, 130.

***Exorides espeletiae* (Kirsch, 1889: 23)**

Distribution: Colombia ('**Bogotá**').

References: Kirsch 1889, 23 (as *Compsus espeletiae*), Marshall 1922b, 217; Wibmer and O'Brien 1986a, 80; Morrone 1999, 130.

***Exorides labyrinthicus* (Kirsch, 1889: 22)**

Distribution: Colombia (**Pasto, Santa Lucía**).

References: (Kirsch 1889, 22 (as *Compsus labyrinthicus*), Marshall 1922b, 203; Wibmer and O'Brien 1986a, 80; Morrone 1999, 130).

***Exorides lindigi* (Kirsch, 1889: 24)**

Distribution: Colombia (**Bogotá**).

References: Kirsch 1889, 24 (as *Compsus lindigi*), Marshall 1922b, 205 (in key), Wibmer and O'Brien 1986a, 80; Morrone 1999, 130.

***Exorides marshalli* (Bovie, 1908b: 44)**

Distribution: **Colombia**.

References: Bovie 1908b, 44 (as *Compsus marshalli*), Wibmer and O'Brien 1986a, 80; Morrone 1999, 130.

***Exorides masoni* Marshall, 1922: 215**

Distribution: Colombia (**Magdalena, San Lorenzo Mt.**).

References: Marshall 1922b, 215; Wibmer and O'Brien 1986a, 80; Morrone 1999, 130.

***Exorides mucronatus* (Faust, 1892: 18)**

Distribution: Colombia, Venezuela.

References: Faust 1892, 18 (as *Synthlibonotus mucronatus*), Marshall 1922b, 214; Wibmer and O'Brien 1986a, 80; Morrone 1999, 130.

***Exorides obesus* Marshall, 1922: 209**

Distribution: **Colombia**.

References: Marshall 1922b, 209; Wibmer and O'Brien 1986a, 80; Morrone 1999, 130.

***Exorides pyriformis* Marshall, 1922: 211**

Distribution: **Colombia**.

References: Marshall 1922b, 211; Wibmer and O'Brien 1986a, 80; Morrone 1999, 130.

***Exorides quadrivittatus* (Kirsch, 1889: 23)**

Distribution: Colombia (**Pasto, Santa Lucía**).

References: Kirsch 1889, 23; Marshall 1922b, 206; Wibmer and O'Brien 1986a, 80; Morrone 1999, 130.

***Exorides rugosus* (Taschenberg, 1870: 188)**

Distribution: Colombia (**Bogotá**).

References: Taschenberg 1870, 188 (as *Compsus rugosus*), Marshall 1922b, 204; Wibmer and O'Brien 1986a, 80; Morrone 1999, 130.

***Exorides septemcostatus* Marshall, 1922: 212**

Distribution: **Colombia**, Ecuador.

References: Marshall 1922b, 212; Wibmer and O'Brien 1986a, 80; Morrone 1999, 130.

Genus *Oxyderces* Schönherr, 1823: col. 1140

Distribution: Argentina, Bolivia, Brazil, Colombia, Ecuador, Paraguay, Peru, Venezuela; Guadeloupe, Martinique.

References: Schönherr 1823, col. 1140; Marshall 1922b, 201 (as *Plococompsus*), O'Brien and Wibmer 1982, 58; Wibmer and O'Brien 1986a, 80; Alonso-Zarazaga and Lyal 1999, 158; Morrone 1999, 130; Girón and Chamorro 2020, 41 (part).

***Oxyderces exaratus* (Hustache, 1938a: 111)**

Distribution: Colombia [Columbia] ([**Transylvanie**] Pensilvania, Cundinamarca (?)).

References: Hustache 1938a, 111 (as *Plococompsus exaratus*), Wibmer and O'Brien 1986a, 81; Morrone 1999, 130.

***Oxyderces mirandus* (Pascoe, 1880: 423)**

Distribution: **Colombia** [Columbia] (Boyacá: Muzo), Ecuador.

References: Pascoe 1880, 423; Marshall 1922b, 201 (listed as *Compsus mirandus*), Hustache 1938a, 114 (as *Plococompsus texatus*), Wibmer and O'Brien 1986a, 81; Morrone 1999, 131.

***Oxyderces viridiaeris* (Hustache, 1938a: 115)**

Distribution: **Colombia**.

References: Hustache 1938a, 115 (as *Plococompsus viridiaeris*), Wibmer and O'Brien 1986a, 81; Morrone 1999, 131.

***Oxyderces viridipes* (Boheman, 1840: 179)**

Distribution: Colombia (**Antioquia**).

References: Boheman in Schönherr 1840b, 179 (as *Platyomus viridipes*), Wibmer and O'Brien 1986a, 81; Morrone 1999, 131.

Genus *Synthlibonotus* Schönherr, 1847: 41 [91]

Distribution: Colombia, Venezuela; Guatemala, Mexico.

References: Schönherr 1847, 41 [91], O'Brien and Wibmer 1982, 58; Wibmer and O'Brien 1986a, 81; Alonso-Zarazaga and Lyal 1999, 158; Morrone 1999, 132.

***Synthlibonotus rufipes* Schönherr, 1847: 41 [91]**

Distribution: **Colombia**, Venezuela.

References: Schönherr 1847, 41 [91], Wibmer and O'Brien 1986a, 81; Morrone 1999, 132.

Genus *Xestogaster* Marshall, 1922: 221

Distribution: Brazil, Colombia, Peru.

References: Marshall 1922b, 221; Wibmer and O'Brien 1986a, 81; Alonso-Zarazaga and Lyal 1999, 158; Morrone 1999, 132.

***Xestogaster porosa* Marshall, 1922: 221**

Distribution: Colombia ('**Bogotá**').

References: Marshall 1922b, 221; Wibmer and O'Brien 1986a, 81; Morrone 1999, 132.

***Xestogaster squalida* Marshall, 1922: 222**

Distribution: **Colombia**.

References: Marshall 1922b, 222; Wibmer and O'Brien 1986a, 81; Morrone 1999, 132.

Tribe Lordopini Schönherr, 1823: col. 1142

Genus *Acanthobrachis* Jekel, 1854 [1857]-I: 9bis (in key) [1857: 101]

Distribution: Brazil, Colombia.

References: Jekel 1857, 101; Kessel 1932, 35; Wibmer and O'Brien 1986a, 91; Alonso-Zarazaga and Lyal 1999, 161; Morrone 1999, 109.

Acanthobrachis germari Jekel, 1857: 103

Distribution: Brazil, Colombia.

References: Jekel 1857, 103; Kessel 1932, 35; Wibmer and O'Brien 1986a, 91; Morrone 1999, 109.

Genus *Atomorhinus* Hustache, 1946: 3

Distribution: Colombia.

References: Hustache 1946, 3; Wibmer and O'Brien 1986a, 94; Alonso-Zarazaga and Lyal 1999, 161; Morrone 1999, 110.

***Atomorhinus impressidorsum* Hustache, 1946: 4**

Distribution: Colombia (Cauca, Rio Chili; Cundinamarca: Fusagasugá).

References: Hustache 1946, 4; Wibmer and O'Brien 1986a, 94; Morrone 1999, 110.

Genus *Deroconus* Jekel, 1854 [1857]-I: 9bis (in key)

Distribution: Colombia, Venezuela.

References: Jekel 1857; Kessel 1932, 62 (as *Deraconus*), Wibmer and O'Brien 1986a, 93; Alonso-Zarazaga and Lyal 1999, 161; Morrone 1999, 111.

Deroconus rufipes (Lacordaire, 1863: 269)

Distribution: **Colombia**, Venezuela.

References: Lacordaire 1863, 269 (as *Hypsonotus rufipes*), Kessel 1932, 66 (as *Deraconus rufipes*), Wibmer and O'Brien 1986a, 93; Morrone 1999, 111.

Genus *Eudmetus* Jekel, 1856: 9bis [1857: 105]

Distribution: Colombia.

References: Jekel 1857, 105; Kessel 1932, 37 (in key), Wibmer and O'Brien 1986a, 91; Alonso-Zarazaga and Lyal 1999, 161; Morrone 1999, 112.

Eudmetus cinereus Jekel, 1857: 107

Distribution: **Colombia**.

References: Jekel 1857, 107; Wibmer and O'Brien 1986a, 91; Morrone 1999, 112.

Eudmetus nigromaculatus Jekel, 1857: 111

Distribution: **Colombia**.

References: Jekel 1856, 111; Wibmer and O'Brien 1986a, 91; Morrone 1999, 112.

Eudmetus posticatus Jekel, 1857: 109

Distribution: **Colombia**.

References: Jekel 1857, 109; Wibmer and O'Brien 1986a, 91; Morrone 1999, 112.

Genus *Granadia* Kessel, 1935: 95

Distribution: Colombia.

References: Kessel 1935, 95 (not seen), Wibmer and O'Brien 1986a, 95; Alonso-Zarazaga and Lyal 1999, 161; Morrone 1999, 113.

***Granadia humeralis* Kessel, 1935: 96**

Distribution: Colombia.

References: Kessel 1935, 96 (not seen), Wibmer and O'Brien 1986a, 95; Morrone 1999, 113.

Genus *Hypoptophila* Voss, 1934: 102

Distribution: Brazil, Colombia.

References: Voss 1934, 102; Wibmer and O'Brien 1986a, 93; Alonso-Zarazaga and Lyal 1999, 161; Morrone 1999, 113.

***Hypoptophila munda* Voss, 1934: 102**

Distribution: **Colombia** [Nov. Granada].

References: Voss 1934, 102; Wibmer and O'Brien 1986a, 93; Morrone 1999, 113.

Genus *Hypoptus* Jekel 1856: 9bis (in key)

Distribution: Bolivia, Colombia, Ecuador; Belize, Costa Rica, El Salvador, Guatemala, Mexico, Nicaragua, Panama; Grenada, St. Vincent.

References: Jekel 1856; Kessel 1932, 57 (incl. key), Wibmer and O'Brien 1986a, 93; Alonso-Zarazaga and Lyal 1999, 161; Morrone 1999, 113.

***Hypoptus arcticus* Kessel, 1932: 62**

Distribution: Colombia ([**Magdalenenstrom**] Magdalena river).

References: Kessel 1932, 62; Wibmer and O'Brien 1986a, 93; Morrone 1999, 113.

***Hypoptus lepyroides* Kessel, 1932: 63**

Distribution: **Colombia** [**Columbien**].

References: Kessel 1932, 63; Wibmer and O'Brien 1986a, 93; Morrone 1999, 113.

Hypoptus macularis Champion, 1911: 303

Distribution: Colombia; Belize, Costa Rica, El Salvador, Guatemala, Mexico, Nicaragua, Panama.

References: Sharp and Champion 1911, 303; Kessel 1932, 61; Wibmer and O'Brien 1986a, 93; Morrone 1999, 113.

Hypoptus setosulus Kessel, 1932: 64

Distribution: **Colombia**.

References: Kessel 1932, 64; Wibmer and O'Brien 1986a, 93; Morrone 1999, 113.

Genus *Hypsonotus* Germar, 1824: 367

Distribution: Argentina, Bolivia, Brazil, Colombia, Ecuador, Paraguay, Venezuela; Costa Rica, Guatemala, Honduras, Mexico; St. Vincent.

References: Germar 1824, 367; Jekel 1854, 137, O'Brien and Wibmer 1982, 66; Wibmer and O'Brien 1986a, 96; Alonso-Zarazaga and Lyal 1999, 162; Morrone 1999, 113.

Hypsonotus acutipennis Jekel, 1859: 229

Distribution: Colombia ('**Bogotá**').

References: Jekel 1856, 160 (listed), 1859, 229 (described), Wibmer and O'Brien 1986a, 97; Morrone 1999, 113.

Hypsonotus apicatus Jekel, 1857: 173

Distribution: Colombia ('**Bogotá**').

References: Jekel 1857, 145 (listed), 1859, 173 (described), Wibmer and O'Brien 1986a, 97; Morrone 1999, 114.

Hypsonotus bipunctatus Jekel, 1859: 233

Distribution: **Colombia**.

References: Jekel 1856, 160 (listed), 1859, 233 (described), Wibmer and O'Brien 1986a, 97; Morrone 1999, 114.

Hypsonotus callosicollis Jekel, 1857: 179

Distribution: Brazil, Colombia.

References: Jekel 1857, 146 (listed), 179 (described), Wibmer and O'Brien 1986a, 97; Morrone 1999, 114.

Hypsonotus compressipennis Jekel, 1859: 235

Distribution: **Colombia**.

References: Jekel 1857, 160 (listed), 1859, 235 (described), Wibmer and O'Brien 1986a, 97; Morrone 1999, 114.

Hypsonotus glaber Kessel, 1937: 170

Distribution: Colombia.

References: Kessel 1937, 170 (not seen), Wibmer and O'Brien 1986a, 98; Morrone 1999, 114.

Hypsonotus hondurensis Kessel, 1937: 161

Distribution: Colombia; Honduras.

References: Kessel 1937, 161 (not seen), O'Brien and Wibmer 1982, 66; Wibmer and O'Brien 1986a, 98; Morrone 1999, 115.

Hypsonotus interior Kessel, 1937: 171

Distribution: Colombia.

References: Kessel 1937, 171 (not seen), Wibmer and O'Brien 1986a, 98; Morrone 1999, 115.

Hypsonotus laevicollis Jekel, 1859: 227

Distribution: Colombia (**Bogotá**).

References: Jekel 1857, 160 (listed), 1859, 227 (described), Wibmer and O'Brien 1986a, 98; Morrone 1999, 115

***Hypsonotus latissimus* Kessel, 1937: 125**

Distribution: Colombia.

References: Kessel 1937, 125 (not seen), Wibmer and O'Brien 1986a, 98; Morrone 1999, 115.

***Hypsonotus limbifer* Kessel, 1937: 155**

Distribution: Colombia.

References: Kessel 1937, 155 (not seen), Wibmer and O'Brien 1986a, 98; Morrone 1999, 115.

***Hypsonotus nitidulus* Jekel, 1859: 237**

Distribution: **Colombia.**

References: Jekel 1857, 160 (listed), 1859, 237 (described), Wibmer and O'Brien 1986a, 98; Morrone 1999, 115.

***Hypsonotus obsoletus* Jekel, 1857: 155**

Distribution: Brazil, **Colombia.**

References: Jekel 1857, 155 (listed), 1859, 227 (described), Wibmer and O'Brien 1986a, 99; Morrone 1999, 115.

***Hypsonotus punctum* Jekel, 1859: 231**

Distribution: **Colombia.**

References: Jekel 1857, 160 (listed), 1859, 231 (described), Wibmer and O'Brien 1986a, 99; Morrone 1999, 116.

***Hypsonotus ramosus* Jekel, 1859: 225**

Distribution: **Colombia.**

References: Jekel 1856, 160 (listed), 1859, 225 (described), Wibmer and O'Brien 1986a, 99; Morrone 1999, 116.

***Hypsonotus setarius* Jekel, 1857: 156**

Distribution: **Colombia.**

References: Jekel 1857, 156; Wibmer and O'Brien 1986a, 99; Morrone 1999, 116.

***Hypsonotus vestitus* Jekel, 1859: 223**

Distribution: **Colombia.**

References: Jekel 1856, 159 (listed), 1859, 223 (described), Wibmer and O'Brien 1986a, 99; Morrone 1999, 116.

Genus *Lordops* Schönherr, 1823: col. 1142

Distribution: Argentina, Brazil, Colombia, Paraguay, Venezuela; Mexico.

References: Schönherr 1823, col. 1142; Jekel 1854, I.II.7, Kessel 1932, 1 (including key to species), O'Brien and Wibmer 1982, 66; Wibmer and O'Brien 1986a, 89; Alonso-Zarazaga and Lyal 1999, 162; Morrone 1999, 116.

Lordops conjugatus Jekel, 1854: I.V.7

Distribution: Brazil, Colombia ('Bogotá'), Paraguay.

References: Jekel 1854, I.V.7, Kessel 1932, 7 (in key; p. 30 comments), Wibmer and O'Brien 1986a, 90; Morrone 1999, 117.

Genus *Orthocnemus* Jekel, 1856: 131

Distribution: Colombia, Ecuador, Peru.

References: Jekel 1856, 131; Kessel 1932, 40; Wibmer and O'Brien 1986a, 92; Alonso-Zarazaga and Lyal 1999, 162; Morrone 1999, 118.

***Orthocnemus lebasii* Jekel, 1857: 133**

Distribution: Colombia (**Bogotá**).

References: Jekel 1857, 133; Kessel 1932, 41; Wibmer and O'Brien 1986a, 92; Morrone 1999, 118.

Genus *Sulla* Kessel, 1937: 182

Distribution: Colombia.

References: Kessel 1937, 182; Wibmer and O'Brien 1986a, 91 (as *Sullana*), Alonso-Zarazaga and Lyal 1999, 162; Morrone 1999, 121 (as *Sullana*).

***Sulla columbiana* Kessel, 1937: 183**

Distribution: Colombia.

References: Kessel 1937, 183; Wibmer and O'Brien 1986a, 91 (as *Sullana columbiana*), Morrone 1999, 121 (as *Sullana columbiana*).

Tribe Naupactini Gistel, 1856: 374

Genus *Amphideritus* Schönherr, 1840: 117

Distribution: Bolivia, Chile, Colombia, Peru.

References: Schönherr 1840b, 117; Wibmer and O'Brien 1986a, 55; Alonso-Zarazaga and Lyal 1999, 163; Morrone 1999, 154 (as synonym of *Macrostylus*), del Río and Lanteri 2011a.

***Amphideritus rugicollis* Kirsch, 1868: 223**

Distribution: Colombia ('Bogotá').

References: Kirsch 1868, 223; Wibmer and O'Brien 1986a, 55; Morrone 1999, 156.

***Amphideritus setosus* (Schönherr, 1847: 29)**

Distribution: **Colombia** (Bogotá).

References: Schönherr 1847, 29 (as *Pterotropis setosus*); Kirsch 1868, 224 (as *Amphideritus squamosus*, a synonym), Wibmer and O'Brien 1986a, 55; Morrone 1999, 156.

***Amphideritus vilis* Boheman, 1840: 118**

Distribution: Colombia (Bogotá).

References: Boheman in Schönherr 1840b, 118; Wibmer and O'Brien 1986a, 55; Morrone 1999, 156.

Genus *Asymmathetes* Wibmer and O'Brien, 1986: 53

Distribution: Colombia, Ecuador.

References: Wibmer and O'Brien 1986a, 53; Alonso-Zarazaga and Lyal 1999, 164; Morrone 1999, 151.

***Asymmathetes steinheili* (Kirsch, 1889: 13)**

Distribution: Colombia (**Consacá**).

References: Kirsch 1889, 13 (as *Naupactus steinheili*), Wibmer and O'Brien 1986a, 53; Morrone 1999, 151.

Genus *Chamaelops* Kirsch, 1868: 235

Distribution: Colombia.

References: Kirsch 1868, 235; Wibmer and O'Brien 1986a, 51; Alonso-Zarazaga and Lyal 1999, 164; Morrone 1999, 152.

***Chamaelops fissirostris* (Chevrolat, 1878: LV)**

Distribution: Colombia (**Bogotá**).

References: Chevrolat 1878, LV (as *Temnoscopus fissirostris*), Wibmer and O'Brien 1986a, 51; Morrone 1999, 151.

***Chamaelops munitus* Kirsch, 1868: 236**

Distribution: Colombia (**Bogotá**).

References: Kirsch 1868, 236; Wibmer and O'Brien 1986a, 51; Morrone 1999, 152.

Genus *Ericydeus* Pascoe, 1880: 422

Distribution: Argentina, Bolivia, Brazil, Colombia, Ecuador, French Guiana, Peru, Venezuela; Costa Rica, Mexico; SW USA.

References: Pascoe 1880, 422; Wibmer and O'Brien 1986a, 51; Lanteri 1995; Alonso-Zarazaga and Lyal 1999, 164; Morrone 1999, 152.

Ericydeus nigropunctatus (Chevrolat, 1877: 170)

Distribution: Colombia (Guainía), Ecuador, Peru, Venezuela.

References: Chevrolat 1877, 170 (as *Cyphus nigropunctatus*), Wibmer and O'Brien 1986a, 51; Lanteri 1995, 402; Morrone 1999, 152.

Ericydeus sedecimpunctatus (Linnaeus, 1758: 386)

Distribution: Argentina, Bolivia, Brazil, Colombia, French Guiana, Guyana, Venezuela; Costa Rica, Panama.

References: Linnaeus 1758, 386 (as *Curculio 16-punctatus*), Wibmer and O'Brien 1986a, 51; Lanteri 1995, 404; Morrone 1999, 153.

Genus *Hoplopactus* Jekel, 1875: 138

Distribution: Bolivia, Brazil, Colombia, Ecuador, French Guiana, Peru, Trinidad, Venezuela.

References: (Jekel 1875, 138; Wibmer and O'Brien 1986a, 54; Alonso-Zarazaga and Lyal 1999, 164; Morrone 1999, 153).

Hoplopactus dentipes (Kirsch, 1868: 231)

Distribution: Colombia ('**Bogotá**').

References: Kirsch 1868, 231 (as *Mimographus dentipes*), Wibmer and O'Brien 1986a, 54; Morrone 1999, 153.

Hoplopactus pavidus (Boheman, 1840: 107)

Distribution: Colombia, Trinidad, Venezuela.

References: Boheman in Schönherr 1840b, 107 (as *Naupactus pavidus*), Wibmer and O'Brien 1986a, 54; Morrone 1999, 154.

Hoplopactus rufipes (Kirsch, 1868: 230)

Distribution: Colombia ('**Bogotá**').

References: Kirsch 1868, 230 (as *Mimographus rufipes*), Wibmer and O'Brien 1986a, 54; Morrone 1999, 154.

***Hoplopactus suturalis* (Kirsch, 1868: 230)**

Distribution: Colombia ('**Bogotá**').

References: Kirsch 1868, 230 (as *Mimographus suturalis*), Wibmer and O'Brien 1986a, 54; Morrone 1999, 154.

Genus *Lanterius* Alonso-Zarazaga y Lyal, 1999

Distribution: Argentina, Brazil, Colombia, Ecuador, Paraguay, Venezuela; Mexico. Panama.

References: Alonso-Zarazaga and Lyal 1999, 164, del Río and Lanteri 2018, 331.

***Lanterius amandus* (Kirsch, 1868: 225)**

Distribution: Colombia ('**Bogotá**'); Panama.

References: Kirsch 1868, 225 (as *Mimographus amandus*), Wibmer and O'Brien 1986a, 56 (as *Mimographus amandus*), Alonso-Zarazaga and Lyal 1999, 164; Morrone 1999, 154 (as *Macrostylus amandus*).

***Lanterius ardosiacus* (Kirsch, 1868: 229), comb. nov.**

Distribution: Colombia ('**Bogotá**').

References: Kirsch 1868, 229 (as *Mimographus ardosiacus*), Wibmer and O'Brien 1986a, 56 (as *Mimographus ardosiacus*), Morrone 1999, 155 (as *Macrostylus ardosiacus*).

***Lanterius cinereoguttatus* (Champion, 1911: 228), comb. nov.**

Distribution: Colombia, Venezuela; Central America (unspecified).

References: Sharp and Champion 1911, 228 (as *Steirarrhinus cinereoguttatus*), Wibmer and O'Brien 1986a, 56 (as *Mimographus cinereoguttatus*), Morrone 1999, 155 (as *Macrostylus cinereoguttatus*).

***Lanterius hirtus* (Voss, 1953: 60), comb. nov.**

Distribution: Colombia ([**West Cordilleren, Rio Aguacatal**]).

References: Voss 1953, 60 (as *Neonanyptactus hirtus*), Wibmer and O'Brien 1986a, 56 (as *Mimographus hirtus*), Morrone 1999, 155 (as *Macrostylus hirtus*).

Lanterius laesicollis (Schönherr, 1847: 35), **comb. nov.**

Distribution: Colombia, Venezuela.

References: Schönherr 1847, 35 (as *Mimographus laesicollis*), Wibmer and O'Brien 1986a, 56 (as *Mimographus laesicollis*), Morrone 1999, 155 (as *Macrostylus laesicollis*).

Lanterius micans (Kirsch, 1868: 229), **comb. nov.**

Distribution: Colombia ('Bogotá').

References: Kirsch 1868, 229 (as *Mimographus micans*), Wibmer and O'Brien 1986a, 56 (as *Mimographus micans*), Morrone 1999, 155 (as *Macrostylus micans*).

Lanterius rotundicollis (Kirsch, 1889: 15), **comb. nov.**

Distribution: Colombia (Popayán).

References: Kirsch 1889, 15 (as *Mimographus rotundicollis*), Wibmer and O'Brien 1986a, 56 (as *Mimographus rotundicollis*), Morrone 1999, 156 (as *Macrostylus rotundicollis*).

Lanterius versicolor (Kirsch, 1889: 15), **comb. nov.**

Distribution: Colombia ('Pasto').

References: Kirsch 1889, 15 (as *Mimographus versicolor*), Wibmer and O'Brien 1986a, 56 (as *Mimographus versicolor*), Morrone 1999, 156 (as *Macrostylus versicolor*).

Lanterius vittatus (Kirsch, 1889: 14), **comb. nov.**

Distribution: Colombia ('Pasto').

References: Kirsch 1889, 14 (as *Mimographus vittatus*), Wibmer and O'Brien 1986a, 56 (as *Mimographus vittatus*), Morrone 1999, 156 (as *Macrostylus vittatus*).

Genus *Leschenius* del Río, Marvaldi y Lanteri, 2012: 55

Distribution: Colombia, Ecuador.

References: del Río et al. 2012, 55.

Leschenius vulcanorum (Kirsch, 1889: 17)

Distribution: Colombia (Nariño), Ecuador.

References: Kirsch 1889, 17 (as *Canephorus vulcanorum*), Wibmer and O'Brien 1986a, 53 (as *Asymmathetes vulcanorum*), Canchala 1992; Morrone 1999, 151 (as *Asymmathetes vulcanorum*), del Río et al. 2012, 62.

Genus *Litostylus* Faust, 1894: 368

Distribution: Argentina, Brazil, Colombia, French Guiana, Peru, Venezuela; Honduras, Nicaragua, Panama; Antigua, Barbados, Dominica, Guadeloupe, Martinique, Montserrat, Saint Barthelemy, Saint Vincent.

References: Faust 1894, 368, O'Brien and Wibmer 1982, 32; Wibmer and O'Brien 1986a, 48; Alonso-Zarazaga and Lyal 1999, 164; Morrone 1999, 154.

Litostylus diadema (Fabricius, 1787: 116)

Distribution: Argentina, Brazil, Colombia, French Guiana, Peru, Venezuela; Honduras, Nicaragua, Panama.

References: Fabricius 1787, 116 (as *Curculio diadema*), O'Brien and Wibmer 1982, 32; Wibmer and O'Brien 1986a, 48; Morrone 1999, 154.

Genus *Melanocyphus* Jekel, 1875: 143

Distribution: Colombia.

References: Jekel 1875, 143; Wibmer and O'Brien 1986a, 54; Alonso-Zarazaga and Lyal 1999, 164; Morrone 1999, 156, del Río and Lanteri 2007a, 2011a, 59 (in key).

Melanocyphus bispinus (Boheman, 1840: 150)

Distribution: Colombia (**Bogotá**).

References: Boheman in Schönherr 1840b, 150 (as *Cyphus bispinus*), Wibmer and O'Brien 1986a, 54; Morrone 1999, 156, del Río and Lanteri 2007a, 131.

Melanocyphus lugubris (Boheman, 1840: 147)

Distribution: Colombia (**Bogotá**).

References: Boheman in Schönherr 1840b, 147 (as *Naupactus lugubris*), Wibmer and O'Brien 1986a, 54; Morrone 1999, 156, del Río and Lanteri 2007a, 130.

Genus *Mimographus* Schönherr, 1847: 34

Distribution: Colombia, Peru; Belize, Costa Rica, Guatemala, Mexico, Nicaragua, Panama.

References: Schönherr 1847, 34; Wibmer and O'Brien 1986a, 56 (as *Steirarrhinus*), Alonso-Zarazaga and Lyal 1999, 165; Morrone 1999, 154 (as *Macrostylus*).

Mimographus decolor (Boheman, 1840: 106)

Distribution: **Colombia**; Central America (unspecified).

References: Boheman in Schönherr 1840b, 106 (as *Naupactus decolor*), Sharp and Champion 1911, 223 (as *Steirarrhinus conicollis*, a synonym), O'Brien and Wibmer 1982, 32 (as *Steirarrhinus conicollis*), Wibmer and O'Brien 1986a, 56 (as *Steirarrhinus decolor*), Morrone 1999, 155 (as *Macrostylus decolor*).

Mimographus depressicollis (Boheman, 1840: 66)

Distribution: Colombia (**Antioquia**).

References: Boheman in Schönherr 1840b, 66 (as *Naupactus depressicollis*), Wibmer and O'Brien 1986a, 56 (as *Steirarrhinus depressicollis*), Morrone 1999, 155 (as *Macrostylus depressicollis*).

Mimographus elegantulus (Boheman, 1840: 64)

Distribution: Colombia (**Antioquia**).

References: Boheman in Schönherr 1840b, 64 (as *Naupactus elegantulus*), Wibmer and O'Brien 1986a, 56 (as *Steirarrhinus elegantulus*), Morrone 1999, 155 (as *Macrostylus elegantulus*).

Mimographus plicaticollis (Hustache, 1938b: 274)

Distribution: Colombia (Cundinamarca (Fusagasugá), Valle del Cauca (Cali), San Antonio, Rio Cauca).

References: Hustache 1938b, 274 (as *Naupactus plicaticollis*), Wibmer and O'Brien 1986a, 56 (as *Steirarrhinus plicaticollis*), Morrone 1999, 155 (as *Macrostylus plicaticollis*).

Mimographus varians (Boheman, 1840: 65)

Distribution: Colombia (**Antioquia**).

References: Boheman in Schönherr 1840b, 65 (as *Naupactus varians*), Wibmer and O'Brien 1986a, 56 (as *Steirarrhinus varians*), Morrone 1999, 155 (as *Macrostylus varians*).

Genus *Naupactus* Dejean, 1821: 94

Distribution: Argentina, Bolivia, Brazil, Chile, Colombia, Ecuador, French Guiana, Guyana, Paraguay, Peru, Surinam, Uruguay, Venezuela; Costa Rica, Guatemala, Honduras, Mexico; SE, SW USA.

References: Dejean 1821, 94; Wibmer and O'Brien 1986a, 57; Lanteri and Morrone 1995; Bordón 1997; Alonso-Zarazaga and Lyal 1999, 165; Morrone 1999, 157; Scatagliini et al. 2005; Rosas et al. 2011; Lanteri and del Río 2016a, 2017, del Río et al. 2018, del Río and Lanteri 2019.

Naupactus instabilis Boheman, 1840: 71

Distribution: **Colombia**, Venezuela.

References: Boheman in Schönherr 1840b, 71; Wibmer and O'Brien 1986a, 60; Morrone 1999, 159.

Naupactus venezolanus Hustache, 1938: 270

Distribution: Colombia, **Venezuela**.

References: Hustache 1938b, 270, O'Brien and Wibmer 1982, 33 (as *Alceis venezolanus*), Wibmer and O'Brien 1986a, 62; Morrone 1999, 162.

Genus *Platyomus* Sahlberg, 1823: 29

Distribution: Argentina, Bolivia, Brazil, Colombia**, Ecuador, Guyana, Paraguay, Peru, Uruguay, Venezuela; Belize, Costa Rica, El Salvador, Guatemala, Honduras. Mexico; SW USA (Texas).

References: Sahlberg 1823, 29; Wibmer and O'Brien 1986a, 49; Alonso-Zarazaga and Lyal 1999, 165; Morrone 1999, 163; Lanteri and del Río 2016a.

Undetermined species

Distribution: Colombia (Amazonas, Antioquia, Meta, Quindío, Risaralda, Valle del Cauca).

References: Girón 2007b; Cardona-Duque et al. 2018, 254–255.

Genus *Plectrophoroides* Wibmer and O'Brien, 1986: 50

Distribution: Argentina, Bolivia, Brazil, Colombia, French Guiana, Guyana, Peru, Surinam, Venezuela.

References: Wibmer and O'Brien 1986a, 50; Alonso-Zarazaga and Lyal 1999, 165; Morrone 1999, 164.

***Plectrophoroides acuminatus* (Chevrolat, 1879: LXXVII)**

Distribution: **Colombia**.

References: Chevrolat 1879, LXXVII (as *Plectrophorus acuminatus*), Wibmer and O'Brien 1986a, 50; Morrone 1999, 164.

***Plectrophoroides albilabris* (Chevrolat, 1879: LXXVIII)**

Distribution: Colombia ('**Bogotá**').

References: Chevrolat 1879, LXXVIII (as *Plectrophorus albilabris*), Wibmer and O'Brien 1986a, 50; Morrone 1999, 164.

***Plectrophoroides bifasciatus* (Chevrolat, 1879: LXXVII)**

Distribution: Bolivia, **Colombia**.

References: Chevrolat 1879, LXXVII (as *Plectrophorus bifasciatus*), Wibmer and O'Brien 1986a, 50; Morrone 1999, 164.

Genus *Tetragonomus* Champion, 1911: 240

Distribution: Colombia; Panama.

References: Sharp and Champion 1911, 240, O'Brien and Wibmer 1982, 32; Wibmer and O'Brien 1986a, 51; Alonso-Zarazaga and Lyal 1999, 165; Morrone 1999, 165.

***Tetragonomus tuberosus* Champion, 1911: 240**

Distribution: Colombia; Panama.

References: Sharp and Champion 1911, 240, O'Brien and Wibmer 1982, 32; Wibmer and O'Brien 1986a, 51; Morrone 1999, 165.

Tribe Premnotrypini Kuschel, 1956: 187

Genus *Premnotrypes* Pierce, 1914: 348

Distribution: Bolivia, Chile, Colombia, Ecuador, Peru, Venezuela.

References: Pierce 1914, 348; Kuschel 1956, 204; Wibmer and O'Brien 1986a, 89; Alonso-Zarazaga and Lyal 1999, 176; Morrone 1999, 107.

Premnotrypes vorax (Hustache, 1933: 377)

Distribution: Colombia ('**Bogotá**').

References: Hustache 1933, 377 (as *Solanophagus vorax*), Kuschel 1956, 209; Wibmer and O'Brien 1986a, 89; Morrone 1999, 107.

Tribe Tanymecini Lacordaire, 1863: 82

Genus *Airosimus* Howden, 1966: 174

Distribution: Bolivia, Brazil, Colombia, Ecuador, Peru, Paraguay, Venezuela; Costa Rica, Mexico, Panama.

References: Howden 1966; Wibmer and O'Brien 1986a, 71; Alonso-Zarazaga and Lyal 1999, 178; Morrone 1999, 143.

Airosimus robustus (Faust, 1892: 4)

Distribution: Colombia, Venezuela.

References: Faust 1892, 4 (as *Menetypus robustus*), Howden 1966, 196; Wibmer and O'Brien 1986a, 71; Morrone 1999, 143.

***Airosimus semirobustus* Howden, 1966: 199**

Distribution: **Colombia.**

References: Howden 1966, 199; Wibmer and O'Brien 1986a, 71; Morrone 1999, 143.

Genus *Hadromeropsis* Pierce, 1913: 400

Distribution: Argentina, Bolivia, Brazil, Colombia, Ecuador, Paraguay, Peru, Uruguay, Venezuela; Costa Rica, Guatemala, Mexico, Panama.

References: Pierce 1913, 400; Howden 1982; Wibmer and O'Brien 1986a, 68; Alonso-Zarazaga and Lyal 1999, 179; Morrone 1999, 143.

Hadromeropsis alacer Howden, 1982: 85

Distribution: **Colombia** (Bogotá, Valle del Cauca), Ecuador.

References: Howden 1982, 85; Wibmer and O'Brien 1986a, 68; Morrone 1999, 143.

Hadromeropsis annae Anderson, 2008: 65

Distribution: Colombia (**Páramo Sumapaz**).

References: Anderson 2008, 65.

Hadromeropsis gemmifera (Boheman, 1845: 418)

Distribution: Colombia (Bogotá, Magdalena, Santander), **Venezuela**; Guatemala, Panama.

References: Boheman in Schönherr 1845, 418 (as *Hadromerus gemmifer*), Howden 1982, 51 (as *Hadromeropsis gemmifer*), Wibmer and O'Brien 1986a, 69; Morrone 1999, 144.

Hadromeropsis impressicollis (Kirsch, 1868: 233)

Distribution: Colombia (**Bogotá**, Valle del Cauca: Socorro, Boyacá: Muzo).

References: Kirsch 1868, 223 (as *Hadromerus impressicollis*), Howden 1982, 94; Wibmer and O'Brien 1986a, 69; Morrone 1999, 144.

Hadromeropsis magica (Pascoe, 1881: 41)

Distribution: Brazil (?), Colombia (Bogotá, Cundinamarca (Fusagasugá)).

References: Pascoe 1881, 41 (as *Naupactus magicus*), Howden 1982, 114 (as *Hadromeropsis magicus*), Wibmer and O'Brien 1986a, 69; Morrone 1999, 144.

Hadromeropsis mandibularis Howden, 1982: 117

Distribution: Colombia (**St. Antonio** (prob. Valle del Cauca)).

References: Howden 1982, 117; Wibmer and O'Brien 1986a, 69; Morrone 1999, 144.

Hadromeropsis meridiana Howden, 1982: 54

Distribution: Brazil, Colombia (Cundinamarca (**Monte Redondo nr. Bogotá**, Bogotá, nr. Girardot, Quetame), Tolima (Ibagué)).

References: Howden 1982, 54 (as *Hadromeropsis meridianus*), Wibmer and O'Brien 1986a, 69; Morrone 1999, 144.

Hadromeropsis nebulicola Howden, 1982: 88

Distribution: Colombia (Magdalena (**San Lorenzo**)).

References: Howden 1982, 88 (as *Hadromeropsis nebolicolus*), Wibmer and O'Brien 1986a, 69; Morrone 1999, 144.

Hadromeropsis pectinata Howden, 1982: 108

Distribution: Bolivia, Colombia, Peru.

References: Howden 1982, 108 (as *Hadromeropsis pectinatus*), Wibmer and O'Brien 1986a, 69; Morrone 1999, 144.

Hadromeropsis silacea Howden, 1982: 90

Distribution: Brazil (?), **Colombia** (Antioquia (Frontino), Cauca).

References: Howden 1982, 90 (as *Hadromeropsis silaceus*), Wibmer and O'Brien 1986a, 69; Morrone 1999, 144.

Hadromeropsis striata Howden, 1982: 126

Distribution: Colombia [**Paso Bella Vista, above Duriamaina**].

References: Howden 1982, 126 (as *Hadromeropsis striatus*), Wibmer and O'Brien 1986a, 69; Morrone 1999, 144.

Genus *Macropterus* Schönherr, 1840: 419

Distribution: Brazil, Colombia, Peru.

References: Schönherr 1840b, 419; Wibmer and O'Brien 1986a, 72; Alonso-Zarazaga and Lyal 1999, 179; Morrone 1999, 145.

Macropterus chlorostomus Boheman, 1840: 424

Distribution: Brazil, Colombia.

References: Boheman in Schönherr 1840b, 424; Wibmer and O'Brien 1986a, 72; Morrone 1999, 145.

Genus *Pandeleteius* Schönherr, 1834: 129

Distribution: Argentina, Bolivia, Brazil, Chile, Colombia, Ecuador, Paraguay, Peru, Venezuela; Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama; Dominica, Grenada, Guadeloupe, Jamaica, Saint Vincent; E Canada, USA.

References: Schönherr 1834, 129; Howden 1959, 1974, 1976, 1986, 1996, 1998, 2001, 2004, 2008, 2011, O'Brien and Wibmer 1982, 48; Wibmer and O'Brien 1986a, 69; Alonso-Zarazaga and Lyal 1999, 179; Morrone 1999, 146; Girón and Howden 2019.

***Pandeleteius admirabilis* Howden, 1976: 192**

Distribution: Colombia (**Cauca**).

References: Howden 1976, 192; Wibmer and O'Brien 1986a, 69; Morrone 1999, 146.

Pandeleteius andeanus Howden, 1976: 155

Distribution: Colombia (Boyacá (Guateque), Cundinamarca (Guasca-Guachetá, Mesitas del Colegio), Norte de Santander (**Chinácota**, Pamplona), Valle del Cauca (Palmira)), Venezuela.

References: Howden 1976, 155; Wibmer and O'Brien 1986a, 69; Morrone 1999, 146.

***Pandeleteius antiochensis* Howden, 1976: 37**

Distribution: Colombia (Antioquia (**Sonsón**)).

References: Howden 1976, 37; Wibmer and O'Brien 1986a, 70; Morrone 1999, 146.

Pandeleteius arcanus Howden, 1876: 60

Distribution: **Colombia** (?), Venezuela.

References: Howden 1976, 60, 2004, 193; Wibmer and O'Brien 1986a, 70; Morrone 1999, 146.

Pandeteius bordoni Howden, 1976: 171

Distribution: Colombia (Norte de Santander (**Pamplona**)).

References: Howden 1976, 171; Wibmer and O'Brien 1986a, 70; Morrone 1999, 146.

Pandeteius campbelli Howden, 1976: 104

Distribution: Colombia (Magdalena (**San Lorenzo**)).

References: Howden 1976, 104; Wibmer and O'Brien 1986a, 70; Morrone 1999, 146.

Pandeteius campestris Howden, 1976: 73

Distribution: Colombia (Atlántico (Puerto Colombia), Bolívar (Cartagena, Turbaco), Cesar (Valledupar), Córdoba (Lorica), Magdalena (**Río Frío**, Sevilla), Sucre (Toluviejo)), Venezuela.

References: Howden 1976, 73, 2004, 207; Wibmer and O'Brien 1986a, 70; Morrone 1999, 146.

Pandeteius carinipenis Howden, 1976: 123

Distribution: Colombia (Valle del Cauca (**Saladito**)).

References: Howden 1976, 123; Wibmer and O'Brien 1986a, 70; Morrone 1999, 146.

Pandeteius chapini Howden, 1976: 68

Distribution: Colombia (Meta (Restrepo, Villavicencio)), Venezuela.

References: Howden 1976, 68, 2004, 203; Wibmer and O'Brien 1986a, 70; Morrone 1999, 146.

Pandeteius clivus Howden, 1976: 146

Distribution: Colombia (Cauca (**Silvia**)).

References: Howden 1976, 146; Wibmer and O'Brien 1986a, 70; Morrone 1999, 146.

***Pandeteius conirostris* Howden, 1976: 101**

Distribution: Colombia (Magdalena (**San Lorenzo**)).

References: Howden 1976, 101; Wibmer and O'Brien 1986a, 70; Morrone 1999, 147.

***Pandeteius dissimilis* Voss, 1939: 337**

Distribution: Colombia (Candelaria (Valle del Cauca (?))); Costa Rica, Honduras, Panama.

References: Voss 1939, 337; Howden 1976, 185; Wibmer and O'Brien 1986a, 70; Morrone 1999, 147.

***Pandeteius eberhardi* Howden, 1976: 135**

Distribution: Colombia (Valle del Cauca: **Pichindé**).

References: Howden 1976, 135; Wibmer and O'Brien 1986a, 70; Morrone 1999, 147.

***Pandeteius excisus* Howden, 1976: 169**

Distribution: Colombia (Cundinamarca (**La Aguadita**, Fusagasugá, Silvania, Tequendama)).

References: Howden 1976, 169; Wibmer and O'Brien 1986a, 70; Morrone 1999, 147.

***Pandeteius flavus* Howden, 1976: 25**

Distribution: Colombia (Valle del Cauca (Cali (Hacienda Arizona, **Río Jamundí**), Río Pance)).

References: Howden 1976, 25; Wibmer and O'Brien 1986a, 70; Morrone 1999, 147.

***Pandeteius giganteus* Howden, 1976: 94**

Distribution: Colombia (Cauca (**Silvia**)).

References: Howden 1976, 94; Wibmer and O'Brien 1986a, 70; Morrone 1999, 147.

Pandeleteius hadromeroides (Kirsch, 1868: 234)

Distribution: Colombia ('**Bogotá**', Cundinamarca (Bogotá, Anolaima, Fusagasugá, Guayabetal, La Aguadita, La Vega, Monte Redondo, Las Tibayas (between Honda and Bogotá), Tena), Quindío (Calarcá)), Venezuela.

References: Kirsch 1868, 234 (as *Menetypus hadromeroides*), Howden 1976, 149; Wibmer and O'Brien 1986a, 70; Morrone 1999, 147; Solano Rojas and Girón 2019.

Pandeleteius hercules Howden, 1976: 109

Distribution: Colombia (Magdalena (**San Lorenzo**)), Venezuela.

References: Howden 1976, 109; Wibmer and O'Brien 1986a, 70; Morrone 1999, 147.

Pandeleteius humboldti Howden, 1976: 45

Distribution: Colombia (Cundinamarca (**Tequendama**, Mosquera, Zipaquirá-Pacho)).

References: Howden 1976, 45; Wibmer and O'Brien 1986a, 70; Morrone 1999, 147.

Pandeleteius minax Dohrn, 1880: 157

Distribution: Colombia ('**Bogotá**').

References: Dohrn 1880, 157; Howden 1976, 78, 2004, 209; Wibmer and O'Brien 1986a, 70; Morrone 1999, 148.

Pandeleteius mirirostris Howden, 1976: 23

Distribution: Colombia (Cundinamarca (Bogotá, **Fusagasugá**, Melgar, Mesitas del Colegio)).

References: Howden 1976, 23; Wibmer and O'Brien 1986a, 70; Morrone 1999, 148.

Pandeleteius modestus (Faust, 1892: 3)

Distribution: Colombia (Norte de Santander (Santiago)), Venezuela.

References: Faust 1892, 3 (as *Menetypus modestus*), Howden 1976, 180, 2004, 227; Wibmer and O'Brien 1986a, 70; Morrone 1999, 148.

Pandeleteius naupactoides (Pascoe, 1881: 38)

Distribution: **Brazil**, Colombia (Cundinamarca).

References: Pascoe 1881, 38 (as *Pandeleitius naupactoides*), Howden 1976, 119; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius nodifer Champion, 1911: 206

Distribution: Colombia (Bolívar (Cartagena), Cesar (Valledupar), Magdalena (Río Frío, Santa Marta, Sevilla)), Venezuela; Nicaragua, El Salvador; Honduras; Jamaica; SE SW USA.

References: Sharp and Champion 1911, 206; Howden 1970, 46, 1976, 184; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius notabilis Howden, 1976: 128

Distribution: Colombia (Cauca (**Silvia**), Galego (Department unknown)).

References: Howden 1976, 128; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius novagranadae Howden, 1976: 80

Distribution: Colombia (Norte de Santander (Chinácota, Pamplona)), **Venezuela**.

References: Howden 1976, 80; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius olympus Howden, 1976: 57

Distribution: Colombia (Cauca (**Silvia**)).

References: Howden 1976, 57; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius peckorum Howden, 1976: 125

Distribution: Colombia (Norte de Santander (**Chinácota**), Venezuela).

References: Howden 1976, 125; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius pilosipectus Howden, 1976: 159

Distribution: Colombia (Cundinamarca (Bogotá, Laguna Ubaque)), **Ecuador**.

References: Howden 1976, 159; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius procollis Howden, 1976: 84

Distribution: Colombia (Norte de Santander (Quebrada Honda)), Venezuela.

References: Howden 1976, 84; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius pygmaeus Howden, 1976: 70

Distribution: Colombia (Quindío (Calarcá), Valle del Cauca (Cali (**Hacienda La Arizona**, Río Jamundí, Río Pance, nr. Pinchindé)).

References: Howden 1976, 70; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius reductus Howden, 1976: 141

Distribution: Colombia (Antioquia (**El Retiro**)).

References: Howden 1976, 141; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius regina Howden, 1976: 177

Distribution: Colombia (Cundinamarca (Bogotá, La Unión-Bogotá), Norte de Santander (**Pamplona**)), Venezuela.

References: Howden 1976, 177; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius santamartae Howden, 1976: 165

Distribution: Colombia (Magdalena (**San Lorenzo**)).

References: Howden 1976, 165; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius scutellatus Howden, 1976: 97

Distribution: Colombia (Cundinamarca (**Bogotá**)).

References: Howden 1976, 97; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius separatus Howden, 1976: 51

Distribution: Colombia (Norte de Santander (**Pamplona**)).

References: Howden 1976, 51; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius subtilis Howden, 1996: 887

Distribution: Colombia (Meta (Restrepo)), Peru, Venezuela.

References: Howden 1996, 887; Morrone 1999, 148.

Pandeleteius summus Howden, 1976: 143

Distribution: Colombia (Cauca (**Silvia**)).

References: Howden 1976, 143; Wibmer and O'Brien 1986a, 71; Morrone 1999, 148.

Pandeleteius tessellatus Howden, 1976: 189

Distribution: **Colombia**.

References: Howden 1976, 189; Wibmer and O'Brien 1986a, 71; Morrone 1999, 149.

Pandeleteius tinctorius Howden, 1976: 116

Distribution: Colombia (Cundinamarca (**Fusagasugá**, Zipaquirá-Pacho)).

References: Howden 1976, 116; Wibmer and O'Brien 1986a, 71; Morrone 1999, 149.

Pandeleteius torquatus Howden, 1976: 62

Distribution: Colombia (Cauca (**Silvia**)).

References: (Howden 1976, 62; Wibmer and O'Brien 1986a, 71; Morrone 1999, 149.

Pandeleteius truncatus Howden, 1976: 138

Distribution: Colombia (Cauca (**Silvia**), Valle del Cauca (Cali)).

References: Howden 1976, 138; Wibmer and O'Brien 1986a, 71; Morrone 1999, 149.

Pandeleteius upsilon Howden, 1976: 90

Distribution: Colombia (Antioquia (**Rionegro**), Valle del Cauca (Lago Calima)).

References: Howden 1976, 90; Wibmer and O'Brien 1986a, 71; Morrone 1999, 149.

Pandeleteius vitticollis Champion, 1911: 202

Distribution: Colombia (Cundinamarca (Arbeláez, Bogotá, Fusagasugá)); Mexico, Guatemala, Honduras, Nicaragua, Panama.

References: Sharp and Champion 1911, 202; Howden 1976, 162; Wibmer and O'Brien 1986a, 71; Morrone 1999, 149.

Additional comments

This paper, as a compilation of information, constitutes the first step towards a better understanding of the biodiversity of entimines as a whole in Colombia and northern South America. It also highlights the areas and groups where a lot of work has been done and groups in need of revisionary work.

Due to its geographic location and attributes, Colombia harbours elements of different faunal components of the Americas: Andean, Amazon, Caribbean, Pacific and Orinoco (Ministerio del Medio Ambiente, Departamento Nacional de Planeación, Instituto Alexander von Humboldt 1996). Colombia is in a key position to contribute to the understanding of distributional, ecological and evolutionary patterns of Entiminae across the Americas, but remains as one of the most prominent knowledge gaps in the region for this subfamily and plenty of other insect taxa.

Except for economically-important species, there is a general lack of knowledge about the natural history of Colombian entimines (e.g. host plants, life cycles, natural enemies), which makes it very difficult to understand some of the patterns observed, for instance, the abundance vs. scarcity of some taxa. On the other hand, most entimines in Colombian collections are vegetation dwellers, whereas leaf litter, a highly diverse microhabitat for weevils (e.g. Anderson 2010), is essentially unexplored.

Most revisionary studies in northern South America have yielded numerous new species (Hustache 1938a; Howden 1976; Bordón 1991, 1997). With the notable exception of Howden (1976), not many taxonomic revisions of Neotropical entimines have included Colombian specimens; many of the specimens studied by Howden were collected directly by her and her colleagues in the early 1970s (Howden 1976). In the few recent revisions including Colombian material, examined specimens are housed in international collections and collected by early researchers in the times of “Nova Granada” (e.g. del Río and Lanteri 2007a, for *Melanocyphus*). This highlights a major issue for advancing the knowledge of Colombian entomofauna, in general: access to Colombian specimens by researchers is greatly limited. This fact, in turn, highlights other problems: (1) type material or even specimens correctly identified to species are extremely scarce (or non-existent) in national collections; (2) national experts (in Curculionidae), who would potentially be able to study type material abroad, are even more scarce, considering the high diversity of the group in the country (see Girón and Cardona-Duque 2018); (3) international experts essentially have no access to recent Colombian specimens, unless they visit Colombian national collections, which may not always be feasible; and, even if international researchers are able to visit, (4) national regulations make it very difficult to borrow specimens to take them out of the

country for study. All of these issues combined, call for creating or strengthening international collaborations and adapting regulations to allow specimen exchange.

The good news is that digitisation efforts in libraries, and both in national and international biological collections, are contributing to improved access to specimens, data and expertise. Platforms, such as Biodiversity Heritage Library (<https://www.biodiversitylibrary.org/>), provide access to publications of original descriptions, museums are digitising their collections (entering data and imaging specimens, including types; for example, The Natural History Museum in London, (<https://data.nhm.ac.uk/search/entomology>) and SiB Colombia (<https://sibcolombia.net/>; <https://colecciones.biodiversidad.co/>) is providing access to specimen data from national collections. In addition, platforms, like iNaturalist (<https://www.inaturalist.org/home>), are continuously generating records for Colombian species. The greatest limitation to learning and understanding Colombian biodiversity is the availability of national experts who are able to study specimens both in national and international collections. Identifying entimines in Colombia is still a challenging task. Some of the specimens recorded here remain identified only to subfamily (Fig. 11) and are presented here in the hope that those can be revised and determined in the future, especially those from scarcely-explored regions.

Even though this work does not include all the entomological collections in the country and many remain to be revised, each collection revised here harbours a very unique diversity. It is evident that there is collecting bias towards the Andean region, with all other regions, but specially Amazon, Caribbean, Orinoco and the Islands (both Caribbean and Pacific), which are potentially as rich in diversity of species, poorly sampled (see maps in Figs 2–11). The distribution of many species within the country is still to be determined: about one third of the species remain simply recorded from “Colombia” and about another third have been described from specimens collected in ‘Bogotá’ of the 1800s to early 1900s. Studying entimines in national collections and collecting them in poorly sampled regions will likely produce new records and taxa new to science all across the country.

Acknowledgements

The production of this contribution started in 2006 as part of my undergraduate research project; at the time, guidance by Nancy Carrejo and support by Carmen Elisa Posso (MUSENUV) and the Grupo de Investigaciones Entomológicas (Departamento de Biología, Facultad de Ciencias, Universidad del Valle, Cali, Colombia) were fundamental for the development of the project. Robert Anderson has been extremely helpful in my career studying broad-nosed weevils: he provided identifications, literature, advice and encouragement during the early development of this work and for that, I will be eternally grateful. I’m in debt to Collections’ curators, managers and assistants who provided access to specimens and on-site equipment: Claudia Alejandra Medina Uribe and Fabio Arturo Gonzalez Alvarado (at IAvH), Dimitri Forero and José Manuel Ramírez Salamanca (at MPUJ). Juliana Cardona-Duque has supported this contribution in many significant ways, providing data from CEUA and MEPB,

georeferencing all entomine records, revising and editing earlier versions of this manuscript and overall keeping me convinced that getting this compilation of information out in the world is valuable. Analía Lanteri has provided invaluable and critical feedback on several versions of this contribution over the years; thanks to her, the content of this document is much more accurate and comprehensive. The late Charles O'Brien and Nico Franz contributed their expertise, access to specimens and literature and encouragement at several points during the development of this work. Juan José Morrone also provided feedback and encouragement as a reviewer of the very first version of this manuscript, back in 2007. Alexander Ortiz facilitated the initial production of line-drawings. Ivon Babativa assisted databasing specimens at IAvH. My main motivation in producing this manuscript is to make information accessible and easy to find. It has taken me years to become familiar with literature about Colombian entomines and nowadays, I have the bibliographic resources, but living overseas, no access to Colombian specimens. I would like young researchers in Colombia to be able to quickly find information and to study entomine specimens right away! Many of the resources referenced here are available via The Biodiversity Heritage Library (<https://www.biodiversitylibrary.org/>), for which I'm also very grateful.

References

- Alcalá P, Alcázar J (1976) Biología y comportamiento de *Premnotrypes suturicallus* Kuschel (Col. Curculionidae). *Revista Peruana de Entomología* 19: 49–52. <https://sisbib.unmsm.edu.pe/BVRevistas/entomologia/v19/pdf/a10v19.pdf>
- Alcázar J, Cisneros F (1998) Taxonomy and bionomics of the Andean Potato Weevil Complex: *Premnotrypes* spp., and related genera. Impact on a changing world: Program Report 1997–98. International Potato Center, 141–151. https://books.google.com/books?id=aWwV1Pv8_cUC&pg=PA141&#v=onepage&q&f=false
- Alonso-Zarazaga MA, Lyal CHC (1999) A world catalogue of families and genera of Curculionoidea (Insecta: Coleoptera) excluding Scolytidae and Platypodidae. Entomopraxis, Barcelona, Spain, 315 pp.
- Anderson RS (1993) Weevils and plants: Phylogenetic versus ecological mediation of evolution of host plant associations in Curculioninae (Coleoptera: Curculionidae). *Memoirs of the Entomological Society of Canada* 125(S165): 197–232. <https://doi.org/10.4039/entm125165197-1>
- Anderson RS (2008) A new species of flightless *Hadromeropsis* from the Colombian páramo (Coleoptera: Curculionidae; Entiminae; Tanymecini). *Zootaxa* 1879(1): 65–68. <https://doi.org/10.11646/zootaxa.1879.1.8>
- Anderson RS (2010) A taxonomic monograph of the Middle American leaf-litter inhabiting weevil genus *Theognete* Champion (Coleoptera: Curculionidae; Molytinae; Lymanini). *Zootaxa* 2458(1): 1–127. <https://doi.org/10.11646/zootaxa.2458.1.1>
- Anderson RS (2018) 2.2.6. Weevil habitat associations and host evolution/coevolution. In: Morphological and molecular perspectives on the phylogeny, evolution, and classification of weevils (Coleoptera: Curculionoidea): Proceedings from the 2016 International Weevil Meeting. *Diversity (Basel)* 10(3): e64. <https://doi.org/10.3390/d10030064>

- Anderson RS, Howden AT (2002) 131 Curculionidae Latreille 1802. American Beetles, Volume II: Polyphaga: Scarabaeoidea through Curculionoidea. CRC Press, Boca Raton, 722–815.
- Anderson RS, Lanteri AA (2000) New genera and species of weevils from the Galapagos Islands, Ecuador, and Cocos Island, Costa Rica (Coleoptera; Curculionidae; Entiminae; Entimini). American Museum Novitates 2000: 1–15. [https://doi.org/10.1206/0003-0082\(2000\)299%3C0001:NGASOW%3E2.0.CO;2](https://doi.org/10.1206/0003-0082(2000)299%3C0001:NGASOW%3E2.0.CO;2)
- Araujo FE, Cavalcante RD, Cavalcante MLS, Melo QMS (1977) *Hypsonotus* sp. (Col. Curculionidae) attacking groundnut (*Arachis hypogaea* L.) in the State of Ceara, Brazil. Fitosanidade 2: 47. <https://agris.fao.org/agris-search/search.do?recordID=BR19780310907>
- Boisduval JBA (1835) Voyage de découvertes de L'Astrolabe exécuté par ordre du Roi, pendant les années 1826-1827-1828-1829, sous le commandement de M.J. Dumont d'Urville. Faune Entomologique de l'Océan Pacifique, avec illustration des insectes nouveaux recueillis pendant le voyage. Deuxième partie. Coléoptères et autres ordres. J. Tatsu, Paris, [VII +] 716 pp. <https://doi.org/10.5962/bhl.title.35819>
- Bordón C (1991) El género *Macrostylus* Boheman (Col. Curc. Brachyderinae, Naupactini) en Venezuela. Acta Biológica Venezuéllica 13(1–2): 1–50.
- Bordón C (1997) El género *Naupactus* Dejean (Coleoptera: Curculionidae) en Venezuela. Acta Biológica Venezuéllica 17(2): 11–51.
- Bouchard P, Bousquet Y, Davies A, Alonso-Zarazaga M, Alonso-Zarazaga M, Lawrence J, Lyal C, Newton A, Reid C, Schmitt M, Slipinski A, Smith A (2011) Family-group names in Coleoptera (Insecta). ZooKeys 88: 1–972. <https://doi.org/10.3897/zookeys.88.807>
- Bovie A (1908a) Coleoptera. Fam. Curculionidae. Subfam. Entiminae. Genera Insectorum, 1–7. [+ 1.] <https://biodiversitylibrary.org/page/33734015>
- Bovie A (1908b) Notes sur les Curculionides. Annales de la Société entomologique de Belgique 52: 43–44. <https://doi.org/10.5962/bhl.part.5001>
- Bruch C (1932) Metamorfosis de *Entimus nobilis* Oliv. (Coleopt., Curculionidae). Revista de Etología 2: 179–185.
- Bustillo AE, Villegas Isaza MC (1986) El picudito del aliso *Bothynodontes* sp. (Coleoptera: Curculionidae) nueva plaga de forestales en Caldas. Apuntes entomológicos 2: 1–3.
- Calvache G (1987) Capture of adults of *Premnotrypes vorax* (Hustache) with adults of the same species as attractant. Revista Colombiana de Entomología 11(2): 9–14. <https://doi.org/10.25100/socolen.v11i2.10258>
- Canchala MB (1992) Estudios biológicos de *Naupactus* sp. Coleoptera: Curculionidae. Nueva plaga en Nariño. Universidad de Nariño, Facultad de Ciencias Agrícolas.
- Cano DM, Serna FJ, Bustillo AE (2002a) Características anatómicas de una nueva especie de *Compsus* (Coleoptera: Curculionidae) plaga de cítricos en Colombia. Revista Colombiana de Entomología 28(1): 33–41. <https://doi.org/10.25100/socolen.v28i1.9626>
- Cano DM, Bustillo AE, Cárdenas R, Orozco L (2002b) Biología y enemigos nativos del picudo de los cítricos *Compsus* n. sp. (Coleoptera: Curculionidae). Revista Colombiana de Entomología 28(1): 43–52. <https://doi.org/10.25100/socolen.v28i1.9627>
- Cardona-Duque J, Bota-Sierra CA, Correa-Carmona Y, Flórez C, Girón JC, Vélez-Bravo A, Wolff M (2018) Estado del conocimiento de insectos en los Bosques Andinos de Antioquia. In: Quintero-Vallejo E, Benavides AM, Moreno N, González-Caro S (Eds) Bosques Andinos, estado actual y retos para su conservación en Antioquia. Fundación

- Jardín Botánico de Medellín Joaquín Antonio Uribe Programa Bosques Andinos (CO-SUDE), Medellín, Colombia, 241–266. <http://www.observatoriobosquesantioquia.org/Libro-Bosques-Andinos>
- Chevrolat LAA (1877) Descriptions de Curculionides nouveaux. *Annales de la Société Entomologique de France* 5(7): 170–173. <https://www.biodiversitylibrary.org/page/8244498>
- Chevrolat LAA (1878) Descriptions de quatre Curculionides nouveaux, dont l'un est le type d'un genre nouveau dans le groupe des *Megalostylus*. *Annales de la Société entomologique de France, Bulletin des Séances*: LIV–LV. <https://biodiversitylibrary.org/page/8233114>
- Chevrolat LAA (1879) Diagnoses de nouvelles espèces de la famille des Curculionides. *Annales de la Société entomologique de France, Bulletin des Séances*: LXXVII–LXXVIII. <https://biodiversitylibrary.org/page/8250753>
- Chevrolat LAA (1881) Fait connaître quatre nouvelles espèces américaines de Curculionites. *Annales de la Société entomologique de France, Bulletin des Séances* 6: XXXIII. <https://gallica.bnf.fr/ark:/12148/bpt6k6335721w/f588.image>
- Cortázar Gómez JE, López-Pazos SA, Cerón J (2012) Determination of Cry toxin activity and identification of an aminopeptidase N receptor-like gene in *Asymmathetes vulcanorum* (Coleoptera: Curculionidae). *Journal of Invertebrate Pathology* 111(1): 94–98. <https://doi.org/10.1016/j.jip.2012.06.003>
- Cortés-Hernández KA, Anderson RS (2019) Two new species of *Isodrusus* Sharp, 1911 (Coleoptera: Curculionidae: Entiminae: Tanymecini). *The Coleopterists Bulletin* 73(4): 940–944. <https://doi.org/10.1649/0010-065X-73.4.940>
- Cortés-Hernández KA, Morrone JJ (2019) A key to the Mexican genera of Tanymecini (Coleoptera: Curculionidae). *Zootaxa* 4615(1): 46–56. <https://doi.org/10.11646/zootaxa.4615.1.2>
- Cortés-Hernández KA, Morrone JJ (2020) Systematic revision of the genus *Isodacrys* Sharp, 1911 (Coleoptera: Curculionidae: Entiminae: Tanymecini). *PeerJ* 8: e10191. <https://doi.org/10.7717/peerj.10191>
- Dejean PFMA (1821) *Catalogue des Coléoptères de la collection de M. le Baron Dejean*. Chez Crevot, Librairie, Paris, 136 pp. <https://doi.org/10.5962/bhl.title.11259>
- del Río MG, Lanteri AA (2007a) Taxonomic revision of *Melanocyphus* Jekel (Coleoptera: Curculionidae). *Studies on Neotropical Fauna and Environment* 42(2): 127–132. <https://doi.org/10.1080/01650520601102567>
- del Río MG, Lanteri AA (2007b) *Thoraconaupactus*, a new Brazilian genus of broad nosed weevil (Coleoptera: Curculionidae: Entiminae) associated with *Leucaena* (Fabaceae). *Entomological News* 118(5): 459–469. [https://doi.org/10.3157/0013-872X\(2007\)118\[459:TANBGO\]2.0.CO;2](https://doi.org/10.3157/0013-872X(2007)118[459:TANBGO]2.0.CO;2)
- del Río MG, Lanteri AA (2011a) *Obrieniolus*, a new monotypic genus of Naupactini (Coleoptera, Curculionidae, Entiminae) from the Peruvian Andes and its phylogenetic placement. *ZooKeys* 102: 51–60. <https://doi.org/10.3897/zookeys.102.1240>
- del Río MG, Lanteri AA (2011b) Taxonomic revision of the genus *Hadropus* Schoenherr (Coleoptera: Curculionidae) with comments on infraspecific variation. *Transactions of the American Entomological Society* 137: 307–315. <https://doi.org/10.3157/061.137.0304>
- del Río MG, Lanteri AA (2012) Redescription of the genus *Curiades* Pascoe 1880 (Coleoptera: Curculionidae: Entiminae: Naupactini), a potential mimic of Mutillidae from Brazil. *Zootaxa* 3570(1): 82–88. <https://doi.org/10.11646/zootaxa.3570.1.6>

- del Río MG, Lanteri AA (2013) Taxonomic revision of the genus *Stenocyphus* Marshall (Coleoptera, Curculionidae) from Brazil. *ZooKeys* 357: 29–43. <https://doi.org/10.3897/zookeys.357.5854>
- del Río MG, Lanteri AA (2018) New synonymies, new combinations and lectotype designations for weevils in the tribe Naupactini (Coleoptera: Curculionidae: Entiminae). *Zootaxa* 4500(3): 329–340. <https://doi.org/10.11646/zootaxa.4500.3.2>
- del Río MG, Lanteri AA (2019) Recognition of species groups of *Naupactus* Dejean (Coleoptera: Curculionidae) from Argentina and neighboring countries. *PeerJ* 6: e6196. <https://doi.org/10.7717/peerj.6196>
- del Río MG, Lanteri AA (2020) A new enigmatic species of broad-nosed weevil endemic to Brazil and its phylogenetic placement within the tribe Naupactini (Coleoptera: Curculionidae: Entiminae). *Papéis Avulsos de Zoologia* 60(special): e202060(s.i.).30. <https://doi.org/10.11606/1807-0205/2020.60.special-issue.30>
- del Río MG, Lanteri AA, Guedes JVC (2006) Taxonomic revision and cladistic analysis of *Teratopactus* Heller (Coleoptera: Curculionidae). *Invertebrate Systematics* 20(5): 585–602. <https://doi.org/10.1071/IS05047>
- del Río MG, Malvardi AE, Lanteri AA (2012) Systematics and cladistics of a new Naupactini genus (Coleoptera: Curculionidae: Entiminae) from the Andes of Colombia and Ecuador. *Zoological Journal of the Linnean Society* 166(1): 54–71. <https://doi.org/10.1111/j.1096-3642.2012.00833.x>
- del Río MG, Morrone JJ, Lanteri AA (2015) Evolutionary biogeography of South American weevils of the tribe Naupactini (Coleoptera: Curculionidae). *Journal of Biogeography* 42(7): 1293–1304. <https://doi.org/10.1111/jbi.12481>
- del Río MG, Rodriguero MS, Confalonieri VA, Lanteri AA (2018) Molecular and morphological phylogenetic analysis of *Naupactus* Dejean (Curculionidae: Entiminae) and allied genera: the dilemma of classification. *Diversity (Basel)* 10(3): e59. <https://doi.org/10.3390/d10030059>
- Dohrn CA (1880) Exotisches. *Entomologische Zeitung* 41: 149–157. <https://biodiversitylibrary.org/page/8989523>
- Dueñas IOT (1989) Ciclo biológico de *Premnotrypes latithorax*, bajo condiciones de laboratorio, en el Cusco. *Revista Peruana de Entomología* 32: 89–92.
- Evenhuis NL (2020) The Insect and Spider Collections of the World Website. <http://hbs.bishopmuseum.org/codens/>
- Fabricius JC (1787) 1 Mantissa insectorum sistens eorum species nuper detectas adiectis characteribus genericis, differentiis specificis, emendationibus, observationibus. Hafniae, [XX +] 348 pp. <https://doi.org/10.5962/bhl.title.11657>
- Faust J (1892) Reise von E. Simon in Venezuela. Curculionidae. Pars prima. *Stettiner Entomologische Zeitung* 53: 1–44. <https://biodiversitylibrary.org/page/9266298>
- Faust J (1893) Neue Helipinen. *Stettiner Entomologische Zeitung* 54: 3–15. <https://www.biodiversitylibrary.org/page/8932093>
- Faust J (1894) Berichtigung. *Stettiner Entomologische Zeitung* 54: 368. <https://babel.hathitrust.org/cgi/pt?id=uiug.30112077164264&view=1up&seq=374>
- Fischer de Waldheim G (1829) *Museum Historiae Naturalis Universitatis Caesareae Mosquensis. Pars II. Insecta. Typis Universitatis Caesareae, Mosquae*, 147 pp.

- Forster JR (1771) *Novae species insectorum: Centuria I*. T. Davies, B. White, 100 pp. <https://doi.org/10.5962/bhl.title.152194>
- Franz NM (2010a) Redescriptions of critical type species in the Eustylini Lacordaire (Coleoptera: Curculionidae: Entiminae). *Journal of Natural History* 44(1–2): 41–80. <https://doi.org/10.1080/00222930903383495>
- Franz NM (2010b) Revision and phylogeny of the Caribbean weevil genus *Apotomoderes* Dejean, 1834 (Coleoptera, Curculionidae, Entiminae). *ZooKeys* 49: 33–75. <https://doi.org/10.3897/zookeys.49.303>
- Franz NM (2011) *Melathra huyenae* Franz, a new genus and new species of Entimine Weevil (Coleoptera: Curculionidae: Entiminae) from Southwestern Hispaniola. *The Coleopterists Bulletin* 65(4): 352–362. <https://doi.org/10.1649/072.065.0406>
- Franz NM (2012) Phylogenetic reassessment of the *Exophthalmus* genus complex (Curculionidae: Entiminae: Eustylini, Geonemini). *Zoological Journal of the Linnean Society* 164(3): 510–557. <https://doi.org/10.1111/j.1096-3642.2011.00774.x>
- Franz NM, Girón JC (2009) *Scelianoma elydimorpha*, a new genus and new species of entimine weevil from southwestern Puerto Rico (Coleoptera: Curculionidae, Entiminae). *Neotropical Entomology* 38(2): 219–230. <https://doi.org/10.1590/S1519-566X2009000200009>
- Gaiger F (2001) Systematic revision and cladistic analysis of the genus *Rhigus* Schoenherr, 1823 (Coleoptera, Curculionidae). *Revista Brasileira de Entomologia* 45: 43–85.
- Gallego JS, Caicedo AM, Carabalí A, Muñoz JE (2012) Comportamiento alimenticio y de oviposición de *Compsus viridivittatus* (Coleoptera: Curculionidae) en especies de cítricos. *Revista Colombiana de Entomología* 38: 191–195. http://www.scielo.org.co/scielo.php?script=sci_abstract&pid=S0120-04882012000200005&lng=en&nrm=iso&tlng=es
- Gemmiger M, von Harold E (1871) *Catalogus coleopterorum hucusque descriptorum synonymicus et systematicus* (Vol. 8). Curculionidae. Gummi, Monachii, 2181–2668. <https://www.biodiversitylibrary.org/page/9649110>
- Germar EF (1817) (Note) in *Miscellen und Correspondenz-Nachrichten*. *Magazine der Entomologie* 2: 339–341. <https://www.biodiversitylibrary.org/page/25412901>
- Germar EF (1824) *Insectorum species novae aut minus cognitae, descriptionibus illustratae* (Vol. 1). Coleoptera. J. C. Hendelii et filii, Halae, [xxiv +] 624 pp. [+2 pl.] <https://doi.org/10.5962/bhl.title.130964>
- Germar EF (1829) Curculionides. *Allgemeine Encyclopädie der Wissenschaften und Künste, Zwanzigster Theil*. Johann Friedrich Oledifch, Leipzig, 356–359. <https://babel.hathitrust.org/cgi/pt?id=umn.31951002385839g&view=1up&seq=78>
- Gillett CPDT, Barr I (2020) First record of the entimine weevil genus *Rhigus* Schoenherr, 1823 (Coleoptera: Curculionidae: Entiminae: Entimini) in Peru: *Rhigus speciosus* (Linnaeus, 1758) in Madre de Dios. *The Coleopterists Bulletin* 74(1): 195–196. <https://doi.org/10.1649/0010-065X-74.1.195>
- Girón JC (2006) A note on Colombian Entiminae. *Curculio* 52: 6–7. https://www.coleopsoc.org/wp-content/uploads/2018/05/curculio_52.pdf
- Girón JC (2007a) Estudio de los Entiminae (Coleoptera: Curculionidae) depositados en el Museo de Entomología de la Universidad del Valle. Undergrad honors thesis abstract. *Boletín del Museo de Entomología de la Universidad del Valle* 8: 1–34. <https://drive.google.com/file/d/1OZMUW9vzydpFcfmudCo2pJWAFKps80dA/view>

- Girón JC (2007b) Estudio de los Entiminae (Coleoptera: Curculionidae) depositados en el Museo de Entomología de la Universidad del Valle. Undergrad honors thesis abstract. Undergrad. Universidad del Valle.
- Girón JC (2020) Listado de las especies de Entiminae (Coleoptera: Curculionidae) de Colombia. Grupo de Coleopterólogos de Colombia. Checklist dataset. <https://doi.org/10.15472/jdwfao> [accessed via GBIF.org on 2020-12-01]
- Girón JC, Cardona-Duque J (2018) Estado del conocimiento de los Curculionidae en Colombia. In: Deloya C, Gasca Álvarez HJ (Eds) Escarabajos del Neotrópico (Insecta: Coleoptera). S y G editores, Mexico, 167–199.
- Girón JC, Chamorro ML (2020) Variability and distribution of the golden-headed weevil *Compsus auricephalus* (Say) (Curculionidae: Entiminae: Eustylini). Biodiversity Data Journal 8: e55474. <https://doi.org/10.3897/BDJ.8.e55474>
- Girón JC, Franz NM (2010) Revision, phylogeny and historical biogeography of the genus *Apodrosus* Marshall, 1922 (Coleoptera: Curculionidae: Entiminae). Insect Systematics & Evolution 41(4): 339–414. <https://doi.org/10.1163/187631210X538799>
- Girón JC, Franz NM (2012) Phylogenetic assessment of the Caribbean weevil genus *Lachnopus* Schoenherr (Coleoptera: Curculionidae: Entiminae). Invertebrate Systematics 26(1): 67–82. <https://doi.org/10.1071/IS11033>
- Girón JC, Howden AT (2019) Five new species of *Pandeteius* Schönherr, 1834 (Coleoptera: Curculionidae: Entiminae: Tanymecini) from South America. The Coleopterists Bulletin 73(4): 831–845. <https://doi.org/10.1649/0010-065X-73.4.831>
- Girón JC, O'Brien CW, Rose-Smyth MC (2018) On the West Indian weevil genus *Lachnopus* Schönherr, 1840 (Coleoptera: Curculionidae: Entiminae): Descriptions of six new species, a proposal for species-groups, and an annotated checklist. Zootaxa 4423(1): 1–85. <https://doi.org/10.11646/zootaxa.4423.1.1>
- Gistel J (1856) Die Mysterien der Europäischen Insectenwelt. Druck und Verlag von Tobias Dannheimer, Kempten, 544 pp. <https://biodiversitylibrary.org/page/42180002>
- Guérin-Ménéville FE (1855) Catalogue des insectes coléoptères recueillis par M. Gaetano Osculati, pendant son exploration de la région équatoriale, sur les bords du Napo et de l'Amazone. Verhandlungen des Zoologisch-Botanischen Vereins in Wien 5: 573–612. <https://biodiversitylibrary.org/page/12049594>
- Guzmán NV, Lanteri AA, Confalonieri VA (2012) Colonization ability of two invasive weevils with different reproductive modes. Evolutionary Ecology 26(6): 1371–1390. <https://doi.org/10.1007/s10682-012-9564-4>
- Háva J, Rukmane A (2018) Short contribution to the genus *Pachyrhynchus* from the Philippines (Coleoptera: Curculionidae: Pachyrhynchini). Acta Biologica Universitatis Daugavpiliensis 18: 203–205. http://sciences.lv/wp-content/uploads/2019/01/H%C3%A1va_218_2.pdf
- Howden AT (1959) A revision of the species of *Pandeteius* Schonherr and *Pandeteinus* Champion of America north of Mexico (Coleoptera: Curculionidae). Proceedings of the California Academy of Sciences 29: 361–421. <https://biodiversitylibrary.org/page/15657419>
- Howden AT (1961) A revision of the genus *Isodacrys* Sharp (Curculionidae, Tanymecini). The Coleopterists Bulletin 15: 75–95. <https://www.jstor.org/stable/3998942>

- Howden AT (1963) A new species of *Isodrusus*, with notes on *Isodrusus debilis* Sharp (Coleoptera: Curculionidae: Tanymecini). The Coleopterists Bulletin 17(2): 43–46. <https://www.jstor.org/stable/3999156>
- Howden AT (1966) *Airosimus*, a new genus of Neotropical Tanymecini (Coleoptera: Curculionidae). Transactions of the American Entomological Society 92: 173–229. <https://www.jstor.org/stable/25077915>
- Howden AT (1969) The genus *Pandeleiteinus* Champion with the description of a new species from Mexico (Curculionidae, Tanymecini). The Coleopterists Bulletin 23: 76–83. <https://www.jstor.org/stable/3999503>
- Howden AT (1970) The Tanymecini of the West Indies (Coleoptera: Curculionidae). Contributions of the American Entomological Institute 5: 1–73.
- Howden AT (1974) The *Pandeleiteius* subgenus *Exmenetypus* Voss in Central America (Coleoptera: Curculionidae, Tanymecini). The Coleopterists Bulletin 28: 7–16. <https://www.jstor.org/stable/3999528>
- Howden AT (1976) *Pandeleiteius* of Venezuela and Colombia (Curculionidae: Brachyderinae: Tanymecini). Memoirs of the American Entomological Institute 24: 1–310.
- Howden AT (1982) Revision of the New World genus *Hadromeropsis* Pierce (Coleoptera, Curculionidae, Tanymecini). Contributions of the American Entomological Institute 19: 1–180.
- Howden AT (1986) The *Pandeleiteius armatus* group, with description of a new species from Mexico (Coleoptera: Curculionidae). The Coleopterists Bulletin 40: 317–323. <https://www.jstor.org/stable/4008295>
- Howden AT (1993) Revision of the Mesoamerican species of *Airosimus* Howden (Coleoptera: Curculionidae). The Coleopterists Bulletin 47: 249–267. <https://www.jstor.org/stable/4009017>
- Howden AT (1996) Neotropical *Pandeleiteius* (Coleoptera: Curculionidae) with irregular elytral striae. The Canadian Entomologist 128(5): 877–955. <https://doi.org/10.4039/Ent128877-5>
- Howden AT (1998) Review of the *Pandeleiteius biseriatus* species-group (Coleoptera: Curculionidae), with description of a new species. The Canadian Entomologist 130(3): 367–375. <https://doi.org/10.4039/Ent130367-3>
- Howden AT (2001) A new species of *Pandeleiteius* from Peru (Coleoptera, Curculionidae). Revue Française d'Entomologie (N. S.) 23: 171–175.
- Howden AT (2004) Review of the genus *Pandeleiteius* (Coleoptera: Curculionidae) of northern South America and the Lesser Antilles. The Canadian Entomologist 136(2): 181–231. <https://doi.org/10.4039/n03-075>
- Howden AT (2008) The species of *Pandeleiteius* Schoenherr of coastal Chile and Peru (Coleoptera, Curculionidae). Zootaxa 1773(1): 55–62. <https://doi.org/10.11646/zootaxa.1773.1.5>
- Howden AT (2011) On some species of *Pandeleiteius* Schoenherr, 1834, in South America south of the tenth parallel (Coleoptera, Curculionidae: Entiminae: Tanymecini). Zootaxa 2977(1): 50–60. <https://doi.org/10.11646/zootaxa.2977.1.2>
- Hustache A (1933) Deux nouveaux curculionides déprédateurs. Bulletin du Muséum National d'Histoire Naturelle 2: 376–380. <https://biodiversitylibrary.org/page/52068095>
- Hustache A (1938a) *Compsus* sud-américains (Coleoptera Curculionidae). Bulletin de la Société Entomologique de Belgique 78: 67–118.

- Hustache A (1938b) Curculionides nouveaux de l'Amérique méridionale, qui se trouvent dans le Deutsches Entomologisches Institut. Deuxième note. Arbeiten über morphologische und taxonomische Entomologie aus Berlin-Dahlem 5: 265–288.
- Hustache A (1946) Deux curculionides nouveaux de l'Amérique du Sud. Bulletin Mensuel de la Société Linneenne de Lyon 15(1): 3–6. <https://doi.org/10.3406/linly.1946.8222>
- Jekel H (1854) Fabricia Entomologica. recueil d'observations nouvelles sur les insectes; monographies, révisions de groupes et de genres, classifications, synonymies et rectifications, descriptions de genres nouveaux et d'espèces nouvelles. <https://doi.org/10.5962/bhl.title.66012>
- Jekel H (1857) Fabricia Entomologica. recueil d'observations nouvelles sur les insectes; monographies, révisions de groupes et de genres, classifications, synonymies et rectifications, descriptions de genres nouveaux et d'espèces nouvelles. Part 1: 97–184. <https://biodiversitylibrary.org/page/42184434>
- Jekel H (1858) Spicilegia Entomologica II: Descriptions of new Curculionidous beetles collected on the voyage of H. M. S. Herald. Annals & Magazine of Natural History 3: 356–351. <https://www.biodiversitylibrary.org/page/18659257>
- Jekel H (1859) Fabricia Entomologica. recueil d'observations nouvelles sur les insectes; monographies, révisions de groupes et de genres, classifications, synonymies et rectifications, descriptions de genres nouveaux et d'espèces nouvelles. Part 1. 3e Livraison: 185–249. <https://biodiversitylibrary.org/page/42184434>
- Jekel H (1875) Coleoptera Jekeliana adjecta eleutheratorum bibliotheca. Énumération systématique et synonymique des coléoptères européens et exotiques composant la collection de Henri Jekel, observations critiques, description d'espèces nouvelles, reproduction et traduction de genres et espèces publiés dans des ouvrages rares français et étrangers de manière à former insensiblement la Bibliothèque du Coleopteriste. Librairie Zoologique de E. Deyrolle fils, Paris. <https://gallica.bnf.fr/ark:/12148/bpt6k884758k/f5.image>
- Kessel F (1932) Fauna Brasiliensis Coleopterologica. Lieferung 1. Friedländer & Sohn, Berlin, 1–32.
- Kessel F (1935) Fauna Brasiliensis Coleopterologica. Lieferung 3. Friedländer & Sohn, Berlin, 73–113.
- Kessel F (1937) Fauna Brasiliensis Coleopterologica. Lieferung 4. Friedländer & Sohn, Berlin, 114–184.
- Kirsch T (1868) Beiträge zur Käferfauna von Bogotá. (Drittes Stück: Brenthiden und adelognathe Curculionen). Berliner Entomologische Zeitschrift 11(3–4): 215–243. <https://biodiversitylibrary.org/page/9215483>
- Kirsch T (1889) Coleopteren gesammelt in den Jahren 1868–1877 auf einer Reise durch Süd Amerika von Alphons Stübel. Abhandlungen und Berichte des Königl. Zoologischen und Anthropologisch-Ethnographischen Museums zu Dresden. 1888–1889: 1–58. [+ 4 pt] <https://www.biodiversitylibrary.org/page/33470151>
- Kuschel G (1955a) *Compsus serrans* n. sp., gorgojo dañino de la caña de azúcar en Venezuela (Aporte 20 de Coleoptera, Curculionidae). Boletín de Entomología Venezolana 11: 1–8.
- Kuschel G (1955b) Nuevas sinonimias y anotaciones sobre Curculionoidea. Revista Chilena de Entomología 4: 261–312. http://www.insectachile.cl/rchen/pdfs/1955v04/Kuschel_1955b.pdf

- Kuschel G (1956) Revision de los Premnotrypini y adiciones a los Bagoini. Boletín del Museo Nacional de Historia Natural 26: 187–235. http://publicaciones.mnhn.gob.cl/668/articles-63959_archivo_01.pdf
- Lacordaire JT (1863) Histoire naturelle des insectes: Genera des Coléoptères ou exposé méthodique et critique de tous les genres proposés jusqu'ici dans cet ordre d'insectes. Tome Sexième. Contenant la famille des Curculionides. Librairie encyclopédique de Roret, Paris, 637 pp. <https://biodiversitylibrary.org/page/9375856>
- Lanteri AA (1980) Contribución al conocimiento del género *Eurymetopus* Schönherr (Coleoptera, Curculionidae). I- Redescrición del género *Eurymetopus* Schönherr y su estudio comparativo con *Floresianus* Hustache y *Floresianellus* Hustache. Revista de la Sociedad Entomológica Argentina 39(3–4): 263–269. <https://www.biotaxa.org/RSEA/article/view/40838>
- Lanteri AA (1982) Estudio taxonómico del género *Wagneriella* Hustache (Coleoptera, Curculionidae). Revista de la Sociedad Entomológica Argentina 41(1–4): 61–76. <https://www.biotaxa.org/RSEA/article/view/40435>
- Lanteri AA (1984) Revisión sistemática del género *Eurymetopus* Schoenherr (Coleoptera, Curculionidae) mediante la aplicación de técnicas numéricas. Revista de la Sociedad Entomológica Argentina 43(1–4): 247–281. <https://www.biotaxa.org/RSEA/article/view/36972>
- Lanteri AA (1985) Revisión de las especies argentinas del género *Macrostylus*, subgénero *Mimographus* Schoenherr (Coleoptera, Curculionidae). CIPFE-CED Orione Contribuciones en Biología 12: 1–6.
- Lanteri AA (1986) Revisión del Género *Asynonychus* Crotch (Coleoptera: Curculionidae). Revista de la Asociacion de Ciencias Naturales del Litoral 17: 161–174. <https://doi.org/10.14409/natura.v2i17.3496>
- Lanteri AA (1989) Estudio sistemático de los géneros *Trichocyphus* Heller y *Mendozella* Hustache (Coleoptera: Curculionidae). Boletín de la Sociedad de Biología de Concepción 60: 139–147. <https://www.biodiversitylibrary.org/page/31654685>
- Lanteri AA (1990a) Revisión sistemática del género *Cyrtomon* Schönherr (Coleoptera, Curculionidae). Revista Brasileira de Entomologia 34: 387–402.
- Lanteri AA (1990b) Revisión sistemática del género *Priocyphus* Hustache 1939 y creación de los géneros *Priocyphopsis* y *Lamprocyphopsis* (Coleoptera, Curculionidae). Revista Brasileira de Entomologia 34: 403–422.
- Lanteri AA (1990c) Revisión sistemática y análisis filogenético de las especies del género *Enoplopactus* Heller 1921 (Coleoptera, Curculionidae). Boletín de la Sociedad de Biología de Concepción 61: 71–92. <https://biodiversitylibrary.org/page/31654879>
- Lanteri AA (1990d) Systematic revision and cladistic analysis of *Phacepholis* Horn. Southwestern Entomologist 15: 179–204. http://agrilife.org/sswe/files/2017/04/SWE_V15.pdf
- Lanteri AA (1992) Systematics, cladistics and biogeography of a new weevil genus, *Galapaganus* (Coleoptera: Curculionidae) from the Galápagos Islands, and coasts of Ecuador and Perú. Transactions of the American Entomological Society 118: 227–267. <https://www.jstor.org/stable/25078560>
- Lanteri AA (1995) Systematic revision of *Ericydeus* Pascoe (Coleoptera: Curculionidae). Entomologica Scandinavica 26(4): 393–424. <https://doi.org/10.1163/187631295X00071>

- Lanteri AA, del Río MG (2003) Revision of the genus *Briarius* [Fischer de Waldheim] (Coleoptera: Curculionidae). *Insect Systematics & Evolution* 34(3): 281–294. <https://doi.org/10.1163/187631203788964755>
- Lanteri AA, del Río MG (2004) Taxonomic revision of *Thoracocyphus* Emden (Coleoptera: Curculionidae). *Insect Systematics & Evolution* 35(4): 449–456. <https://doi.org/10.1163/187631204788912454>
- Lanteri AA, del Río MG (2005) Taxonomy of the monotypic genus *Trichaptus* Pascoe (Coleoptera: Curculionidae: Entiminae), a potential weevil mimic of Mutillidae. *The Coleopterists Bulletin* 59(1): 47–54. <https://doi.org/10.1649/683>
- Lanteri AA, del Río MG (2006a) Taxonomic revision of the genus *Cyphopsis* Roelofs (Coleoptera, Curculionidae). *Mitteilungen aus dem Museum für Naturkunde in Berlin. Deutsche Entomologische Zeitschrift* 53: 275–281. <https://doi.org/10.1002/mmnd.200600025>
- Lanteri AA, del Río MG (2006b) Taxonomic revision of the monotypic genus *Acyphus* Heller (Coleoptera: Curculionidae) with comments on infraspecific variation. *Zootaxa* 1312(1): 59–68. <https://doi.org/10.11646/zootaxa.1312.1.5>
- Lanteri AA, del Río MG (2008) Caracteres genitales de la hembra en la clasificación y filogenia de la tribu Naupactini (Coleoptera: Curculionidae). In: *Contribuciones taxonómicas en órdenes de insectos hiperdiversos*. UNAM-RIBES-CYTED, 159–176.
- Lanteri AA, del Río MG (2016a) Phylogeny of the tribe Naupactini (Coleoptera: Curculionidae) based on morphological characters. *Systematic Entomology* 42(2): 429–447. <https://doi.org/10.1111/syen.12223>
- Lanteri AA, del Río MG (2016b) Taxonomy and cladistics of the group of genera related to *Cyrtomon* Schoenherr (Coleoptera: Curculionidae: Naupactini). *Revista de la Sociedad Entomológica Argentina* 75: 387–402. <https://www.biotaxa.org/RSEA/article/view/22298>
- Lanteri AA, del Río MG (2017) *Naupactus xanthographus* (Germar) species group (Curculionidae: Entiminae: Naupactini): a comprehensive taxonomic treatment. *Journal of Natural History* 51(27–28): 1557–1587. <https://doi.org/10.1080/00222933.2017.1346715>
- Lanteri AA, Díaz NB (1994) Systematic study and cladistic analysis of the genus *Aramigus* Horn (Coleoptera: Curculionidae). *Transactions of the American Entomological Society* 120: 113–144. <https://www.jstor.org/stable/40785183>
- Lanteri AA, Loíacono MS (1990) Systematic study of the *Pantomorus viridisquamosus* species group (Coleoptera: Curculionidae). *Insecta Mundi* 4: 1–10. <https://digitalcommons.unl.edu/insectamundi/396>
- Lanteri AA, Marvaldi AE (1995) *Graphognathus* Buchanan a new synonym of *Naupactus* Dejean and systematics of the *N. leucoloma* species group (Coleoptera: Curculionidae). *The The Coleopterists Bulletin* 49: 206–228. <https://www.jstor.org/stable/4009138>
- Lanteri AA, Morrone JJ (1991) Cladistic analysis of *Priocyphus* Hustache and related genera (Coleoptera: Curculionidae). *Proceedings of the Entomological Society of Washington* 93: 278–287. <https://biodiversitylibrary.org/page/16197951>
- Lanteri AA, Morrone JJ (1995) Cladistics of the *Naupactus leucoloma* species group, *Atrichonotus*, and *Eurymetopus* (Coleoptera: Curculionidae). *Revista de la Sociedad Entomológica Argentina* 54: 99–112. <https://www.biotaxa.org/RSEA/article/view/33786>

- Lanteri AA, Normark BB (1995) Parthenogenesis in the tribe Naupactini (Coleoptera: Curculionidae). *Annals of the Entomological Society of America* 88(6): 722–731. <https://doi.org/10.1093/aesa/88.6.722>
- Lanteri AA, O'Brien CW (1990) Taxonomic revision and cladistic analysis of *Atrichonotus* Buchanan (Coleoptera: Curculionidae). *Transactions of the American Entomological Society* 116: 697–725. <https://www.jstor.org/stable/25078526>
- Lanteri AA, Guedes JC, Parra JRP (2002) Weevils injurious for roots of *Citrus* in São Paulo state, Brazil. *Neotropical Entomology* 31(4): 561–569. <https://doi.org/10.1590/S1519-566X2002000400008>
- Lanteri AA, Guzmán NV, del Río MG, Confalonieri VA (2013) Potential geographic distributions and successful invasions of parthenogenetic broad-nosed weevils (Coleoptera: Curculionidae) native to South America. *Environmental Entomology* 42(4): 677–687. <https://doi.org/10.1603/EN12297>
- Lanteri AA, Loíacono MS, Coscarón M del C, Díaz NB (2017) Systematic study of the *Pantomorus auripes* species group (Coleoptera: Curculionidae). *Revista de la Sociedad Entomológica Argentina* 49: 3–16. <https://www.biotaxa.org/RSEA/article/view/33985>
- Lázaro HO, Nasca AJ, Frascarolo D, Lemme MC, Labatte de Rudelli ME (1997) *Promecops claviger* (Coleoptera: Curculionidae): Nueva plaga de la soja en Tucumán, Argentina. *Revista de la Sociedad Entomológica Argentina* 56: 167–168. <https://www.biotaxa.org/RSEA/article/view/33683>
- Lázaro HO, Cruz D, Nasca AJ (1998) *Promecops claviger* Hustache (Coleoptera: Curculionidae): aspectos biológicos y daños al cultivo de la soja en Tucumán, R. Argentina. *Acta Entomológica Chilena*: 45–47. <https://dialnet.unirioja.es/servlet/articulo?codigo=2013619>
- LeConte JL (1874) The Classification of the Rhynchophorous Coleoptera. *American Naturalist* 8(8): 452–470. <https://doi.org/10.1086/271360>
- LeConte JL, Horn GH (1876) The Rhynchophora of America, North of Mexico. *Proceedings of the American Philosophical Society* 15(96): [vii–ix + xi–xvi+] 1–442. <https://www.jstor.org/stable/982412>
- Linnaeus C von (1758) *Systema Naturae per Regna Tria Naturae: Secundum Classes, Ordines, Genera, Species, Cum Characteribus, Differentiis, Synonymis, Locis*. Impensis Direct (Vol. 1). Laurentii Salvii, Holmiae, 824 pp. <https://biodiversitylibrary.org/page/726886>
- Lona C (1938) Curculionidae: Otiorrhynchinae III. *Coleopterorum Catalogus*. Dr. W. Junk Verlag für Naturwissenschaften, 's-Gravenhage, 415–600.
- Marshall GAK (1916) On new Neotropical Curculionidae. *Annals & Magazine of Natural History* 8(108): 449–469. <https://doi.org/10.1080/00222931609486900>
- Marshall GAK (1922a) Some injurious Neotropical weevils (Curculionidae). *Bulletin of Entomological Research* 13(1): 59–71. <https://doi.org/10.1017/S0007485300045247>
- Marshall GAK (1922b) VI. On new genera and species of Neotropical Curculionidae. *Transactions of the Royal Entomological Society of London* 1922–1923: 181–224. <https://doi.org/10.1111/j.1365-2311.1922.tb02830.x> [+ pl. III-I]
- Marshall GAK (1926) LXVI – On new Neotropical Curculionidae (Col.). *Annals & Magazine of Natural History* 9(107): 530–543. <https://doi.org/10.1080/00222932608633549>

- Marvaldi AE (1997) Higher level phylogeny of Curculionidae (Coleoptera: Curculionoidea) based mainly on larval characters, with special reference to broad-nosed weevils. *Cladistics* 13(4): 285–312. <https://doi.org/10.1111/j.1096-0031.1997.tb00321.x>
- Marvaldi AE (1998) Larvae of Entiminae (Coleoptera: Curculionidae): tribal diagnoses and phylogenetic key, with a proposal about natural groups within Entimini. *Insect Systematics & Evolution* 29(1): 89–98. <https://doi.org/10.1163/187631298X00212>
- Marvaldi AE, Lanteri AA (2005) Key to higher taxa of South American weevils based on adult characters (Coleoptera, Curculionoidea). *Revista Chilena de Historia Natural* 78(1): 65–87. <https://doi.org/10.4067/S0716-078X2005000100006>
- Marvaldi AE, Lanteri AA, del Río MG, Oberprieler RG (2014) Entiminae Schoenherr, 1823. In: Leschen RAB, Beutel RG (Eds) *Handbook of Zoology, Arthropoda: Insecta: Coleoptera, Volume 3: Morphology and Systematics (Phytophaga)*. De Gruyter, Berlin, 503–522.
- Marvaldi AE, del Río MG, Pereyra VA, Rocamundi N, Lanteri AA (2018) A combined molecular and morphological approach to explore the higher phylogeny of entimine weevils (Coleoptera: Curculionidae), with special reference to South American taxa. *Diversity (Basel)* 10(3): e95. <https://doi.org/10.3390/d10030095>
- Mazo-Vargas A (2011) Untangling the effects of biogeography and host plant associations: phylogenetic and phylogeographic studies in the *Exophthalmus* genus complex. Master of Science Thesis. University of Puerto Rico, at Mayaguez.
- Ministerio del Medio Ambiente, Departamento Nacional de Planeación, Instituto Alexander von Humboldt (1996) Política Nacional de Biodiversidad. https://www.minambiente.gov.co/images/BosquesBiodiversidadyServiciosEcosistemas/pdf/Politica-Nacional-de-Biodiversidad/politica_nacional-biodiversidad.pdf
- Morrone JJ (1999) The species of Entiminae (Coleoptera: Curculionidae) ranged in America South of the United States. *Anales del Instituto de Biología. Serie Zoología* 70(2): 99–168. <http://revistas.unam.mx/index.php/zoo/article/view/7224>
- Morrone JJ (2002) The Neotropical weevil genus *Entimus* (Coleoptera: Curculionidae: Entiminae): cladistics, biogeography, and modes of speciation. *The Coleopterists Bulletin* 56(4): 501–513. [https://doi.org/10.1649/0010-065X\(2002\)056\[0501:TNWGEC\]2.0.CO;2](https://doi.org/10.1649/0010-065X(2002)056[0501:TNWGEC]2.0.CO;2)
- Morrone JJ, Abadie EI, Godinho Jr C (2019) A new species of *Entimus* Germar (Coleoptera: Curculionidae: Entiminae) from southeastern Brazil. *Zootaxa* 4590(1): 191–195. <https://doi.org/10.11646/zootaxa.4590.1.10>
- O'Brien CW, Kovarik PW (2001) The genus *Diaprepes*: its origin and geographical distribution in the Caribbean region. https://irrec.ifas.ufl.edu/flcitrus/short_course_and_workshop/diaprepes/Genus_diaprepes.shtml
- O'Brien CW, Peña J (2012) Two species of *Compsus* Schoenherr, new citrus pests from Colombia (Coleoptera: Curculionidae: Entiminae). *Insecta Mundi* 0227: 1–13. <https://digitalcommons.unl.edu/insectamundi/736>
- O'Brien CW, Wibmer GJ (1982) Annotated checklist of the weevils (Curculionidae sensu lato) of North America, Central America, and the West Indies (Coleoptera: Curculionoidea). *Memoirs of the American Entomological Institute* 34: 1–382.
- Oberprieler RG, Marvaldi AE, Anderson RS (2007) Weevils, weevils, weevils everywhere. *Zootaxa* 1668(1): 491–520. <https://doi.org/10.11646/zootaxa.1668.1.24>

- Olivier GA (1790) Encyclopédie méthodique. Histoire naturelle. Insectes. Volume 5. Chez Panckoucke, Librairie, Paris, 793 pp. <https://www.biodiversitylibrary.org/page/7577222>
- Olivier GA (1807) Entomologie, ou Histoire naturelle des insectes: avec leurs caractères génériques et spécifiques, leur description, leur synonymie, et leur figure enluminée. Coléoptères. Tome Cinquieme. De l'Imprimerie de Baudoin, Paris, 612 pp. <https://doi.org/10.5962/bhl.title.61905>
- Pascoe FP (1872) Contributions towards a knowledge of the Curculionidae. Part III. Zoological Journal of the Linnean Society 11(55): 440–492. <https://doi.org/10.1111/j.1096-3642.1872.tb01668.x>
- Pascoe FP (1880) New Neotropical Curculionidae. Part I. Annals and Magazine of Natural History 5(29): 419–428. <https://doi.org/10.1080/00222938009459433>
- Pascoe FP (1881) VI– New neotropical Curculionidae. Part IV. Annals and Magazine of Natural History. Series 5. 7(37): 38–45. <https://doi.org/10.1080/00222938109459470>
- Peña LA (2001) Gusanos blancos de la papa, biología y manejo. Innovación y cambio tecnológico 2: 29–33.
- Perty M (1832) Insecta Brasiliensia. Delectus animalium articulorum, quae in itinere per Brasiliam, annis MDCCCXVII–MDCCCXX: jussu et auspiciis Maximiliani Josephi I. Impensis Editoris, Monachii, 190 pp. <https://www.biodiversitylibrary.org/page/57932830>
- Pierce WD (1913) Miscellaneous contributions to the knowledge of the weevils of the families Attelabidae and Brachyrhinidae. Proceedings of the United States National Museum 45: 365–426. <https://doi.org/10.5479/si.00963801.1988.365>
- Pierce WD (1914) New potato weevils from Andean South America. Journal of Agricultural Research 1: 347–352. [+ pl. XXXDC–XLI] <https://naldc.nal.usda.gov/download/IND43965426/PDF>
- Poinar Jr G, Brown AE (2011) Descriptions of a broad-nosed weevil (Eudiagogini: Curculionidae) and false ladybird beetle (Nilionini: Nilionidae) in Dominican amber. Historical Biology 23(2–3): 231–235. <https://doi.org/10.1080/08912963.2010.527159>
- Poinar G, Legalov AA (2017) Five new species from the subfamily Entiminae (Coleoptera: Curculionidae) in Dominican amber. Palaeontologia Electronica 20: 1–13. <https://doi.org/10.26879/698>
- Poinar G, Bukejs A, Legalov AA (2017) First records of weevils (Coleoptera: Curculionidae) in Quaternary Colombian copal. Revista Colombiana de Entomologia 43(1): 85–90. <https://doi.org/10.25100/socolen.v43i1.6654>
- R Core Team (2018) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. <https://www.R-project.org/>
- Regueira JCS, Damasceno EM, Iannuzzi L (2020) Shape variation of *Cydianerus latruncularius* (Coleoptera, Curculionidae) across biomes and sexes. Zoologischer Anzeiger 289: 96–107. <https://doi.org/10.1016/j.jcz.2020.10.002>
- Rocha Barreto M, Cavalet F (2016) *Promecops claviger* Hustache, 1929 (Coleoptera: Curculionidae): first register in Brazil and first host plant register. Check List 7(5): 616–620. <https://doi.org/10.15560/7.5.616>
- Romo A, Morrone JJ (2011) Track analysis of the Neotropical Entimini (Coleoptera: Curculionidae: Entiminae). Revista Brasileira de Entomologia 55(3): 313–316. <https://doi.org/10.1590/S0085-56262011005000034>

- Rosas MV, Morrone JJ, del Río MG, Lanteri AA (2011) Phylogenetic analysis of the *Pantomorus-Naupactus* complex (Coleoptera: Curculionidae: Entiminae) from North and Central America. *Zootaxa* 2780(1): 1–19. <https://doi.org/10.11646/zootaxa.2780.1.1>
- RStudio Team (2016) RStudio: Integrated Development for R. RStudio, Inc., Boston. <http://www.rstudio.com/>
- Sahlberg CR (1823) Periculi entomographici, species insectorum nondum descriptas proposituri, fasciculus. Frenckel, Aboae, 82 pp. <https://www.biodiversitylibrary.org/page/10990686>
- Say T (1824) Descriptions of coleopterous insects collected in the late expedition to the Rocky Mountains, performed by order of Mr. Calhoun, Secretary of War, under the command of Major Long. By Thomas Say, Zoologist to the Expedition. *Journal of the Academy of Natural Sciences of Philadelphia* 3: 139–216. <https://www.biodiversitylibrary.org/page/24668950>
- Scataglieni MA, Lanteri AA, Confalonieri VA (2005) Phylogeny of the *Pantomorus-Naupactus* complex based on morphological and molecular data (Coleoptera: Curculionidae). *Cladistics* 21(2): 131–142. <https://doi.org/10.1111/j.1096-0031.2005.00055.x>
- Schönherr CJ (1823) Curculionides [Tabula synoptica familiae Curculionidum]. *Isis von Oken* 1823: 1132–1146. <https://biodiversitylibrary.org/page/13257284>
- Schönherr CJ (1834) Genera et species Curculionidum, cum synonymia hujus familiae. Species novae aut hactenus minus cognitae, descriptionibus a Dom. Leonardo Gyllenhal, C. H. Boheman, et entomologis aliis illustratae, tomus secundus, pars prima. Roret, Paris, 673 pp. <https://doi.org/10.5962/bhl.title.8952>
- Schönherr CJ (1840a) Genera et species Curculionidum, cum synonymia hujus familiae. Species novae aut hactenus minus cognitae, descriptionibus a Dom. Leonardo Gyllenhal, C. H. Boheman, et entomologis aliis illustratae, tomus quintus, pars prima. Roret, Paris, 974 pp. <https://www.biodiversitylibrary.org/page/4110075>
- Schönherr CJ (1840b) Genera et species Curculionidum, cum synonymia hujus familiae. Species novae aut hactenus minus cognitae, descriptionibus a Dom. Leonardo Gyllenhal, C. H. Boheman, et entomologis aliis illustratae, tomus sextus, pars prima. Roret, Paris, 474 pp. <https://www.biodiversitylibrary.org/page/4114879>
- Schönherr CJ (1842) Genera et species Curculionidum, cum synonymia hujus familiae. Species novae aut hactenus minus cognitae, descriptionibus a Dom. Leonardo Gyllenhal, C. H. Boheman, et entomologis aliis illustratae, tomus sextus, pars secunda. Roret, Paris, 495 pp. <https://www.biodiversitylibrary.org/page/4115130>
- Schönherr CJ (1845) Genera et species Curculionidum, cum synonymia hujus familiae. Species novae aut hactenus minus cognitae, descriptionibus a Dom. Leonardo Gyllenhal, C. H. Boheman, et entomologis aliis illustratae, tomus octavus, pars secunda. Roret, Paris, 504 pp. <https://www.biodiversitylibrary.org/page/4114205>
- Schönherr CJ (1847) Mantissa secunda familiae curculionidum. Seu descriptiones novorum quorundam generum curculionidum. Typis Norstedt et filii, Holmiae, 86 pp. <https://doi.org/10.5962/bhl.title.5794>
- Scudder SH (1876) Fossil Coleoptera from the Rocky Mountain Tertiaries. *Bulletin of the United States Geological and Geographical Survey of the Territories* 2: 77–87. <https://www.biodiversitylibrary.org/page/32141159>

- Sharp D, Champion GC (1911) *Biologia Centrali-Americana*. Insecta. Coleoptera. Rhynchophora. Volume 4, part 3. R. H. Porter, London, 354 pp. [+ pl. 15] <https://biodiversitylibrary.org/page/581470>
- Shorthouse DP (2010) SimpleMappr, an online tool to produce publication-quality point maps. <https://www.simplemappr.net>
- Solano Rojas Y, Girón JC (2019) First record of *Pandeleiteius hadromeroides* (Kirsch) (Coleoptera: Curculionidae: Entiminae: Tanymericini) on strawberry crops in Venezuela. *The The Coleopterists Bulletin* 73(4): 962–965. <https://doi.org/10.1649/0010-065X-73.4.962>
- Soto-Hernández M, Barros-Barrios MM (2018) Curculiónidos (Coleoptera: Curculionidae) en el cultivo de Nogal pecanero en el estado de Coahuila, México. *Entomología Mexicana* 5: 580–586. <http://www.entomologia.socmexent.org/revista/2018/SM/SM%20580-586.pdf>
- Taschenberg EL (1870) Neue Käfer aus Colombien und Ecuador. *Zeitschrift für die gesammten Naturwissenschaft* 35: 177–199. <https://babel.hathitrust.org/cgi/pt?id=hvd.32044106244114&view=1up&seq=191>
- Tennekes M (2018) tmap: Thematic Maps in R. *Journal of Statistical Software* 84(6): 1–39. <https://doi.org/10.18637/jss.v084.i06>
- Thompson RT (1992) Observations on the morphology and classification of weevils (Coleoptera, Curculionoidea) with a key to major groups. *Journal of Natural History* 26(4): 835–891. <https://doi.org/10.1080/00222939200770511>
- Van Dam MH, Lam AW, Sagata K, Gewa B, Laufa R, Balke M, Faircloth BC, Riedel A (2017) Ultraconserved elements (UCEs) resolve the phylogeny of Australasian smurf-weevils. *PLoS One* 12(11): e0188044. <https://doi.org/10.1371/journal.pone.0188044>
- van Emden FI (1944a) L–A key to the genera of Brachyderinae of the World. *Annals & Magazine of Natural History* 11(81): 559–586. <https://doi.org/10.1080/00222934408527456>
- van Emden FI (1944b) XLVII–A key to the genera of Brachyderinae of the World. *Annals & Magazine of Natural History* 11(80): 503–532. <https://doi.org/10.1080/00222934408527452>
- Vanin SA (1983) Revision of *Phaedropus* Schoenherr with notes on the male genitalia of Entimini (Coleoptera: Curculionidae: Leptopiinae). *The The Coleopterists Bulletin* 37: 283–298. <https://www.jstor.org/stable/4008033>
- Vanin SA, Gaiger F (2005) A cladistic analysis of the genera of the tribe Entimini (Coleoptera, Curculionidae), with description of a new genus and species from the Amazonian Region. *Zootaxa* 1053(1): 1–21. <https://doi.org/10.11646/zootaxa.1053.1.1>
- Vaurie P (1951) Revision of the genus *Entimus* with notes on other genera of Entimini (Coleoptera Curculionidae). *Revista Chilena de Entomología* 1: 147–171. <https://biodiversitylibrary.org/page/32160206>
- Vaurie P (1963) A revision of the South American genus *Hyphantus* (Coleoptera, Curculionidae, Otiorhynchinae). *Bulletin of the American Museum of Natural History* 125: 239–304. <http://digitallibrary.amnh.org/handle/2246/1979>
- Voss E (1934) Einige unbeschriebene neotropische Curculioniden nebst einer Vorstudie zur tribus Promecopini (Col. Curc.). *Sborník Entomologického Oddelení Národního Muzea v Praze* 12: 63–104. http://aemnp.eu/PDF/12_0/12_0_63.pdf

- Voss E (1936) Über einige exotische Curculioniden (Col., Curc). Bulletin de la Société des naturalistes luxembourgeois, N. S. 46: 103–124. https://www.snl.lu/publications/bulletin/SNL_1936_046_103_124.pdf
- Voss E (1939) Über neue und wenig bekannte Rüssler-Arten aus der neotropischen Region, vorwiegend aus Brasilien (Coleoptera: Curculionidae). Arbeiten über morphologische und taxonomische entomologie aus Berlin-Dahlem 6: 335–340.
- Voss E (1953) Neue und bemerkenswerte Curculioniden aus Colombien und Bolivien. (Col. Curc.). Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg 1: 55–84. https://www.zobodat.at/pdf/Ent-Mitt-Zool-Mus-Hamburg_1_0055-0084.pdf
- Warner RE (1979) The genus *Eudiagogus* (Coleoptera, Curculionidae, Leptopiinae), with two new species on the weed *Sesbania* (Leguminosae). Proceedings of the Entomological Society of Washington 81: 304–320. <https://www.biodiversitylibrary.org/part/63464>
- Waterhouse CO (1879) Descriptions of new Coleoptera from Medellin, Colombia, recently added to the British Museum collection. Cistula Entomologica 2: 421–429. <https://biodiversitylibrary.org/page/10891087>
- Wibmer GJ, O'Brien CW (1986a) Annotated checklist of the weevils (Curculionidae sensu lato) of South America (Coleoptera: Curculionoidea). Memoirs of the American Entomological Institute 39: 1–563.
- Wibmer GJ, O'Brien CW (1986b) *Geonemus* Schoenherr, 1833 and *Brachyomus* Lacordaire, 1863 (Insecta, Coleoptera): Proposal to maintain current usage by designation of a type species for *Geonemus*. ZN (S.) 2565. Bulletin of Zoological Nomenclature 43: 300–302. <https://doi.org/10.5962/bhl.part.454>
- Wilts BD, Michielsen K, Kuipers J, De Raedt H, Stavenga DG (2012) Brilliant camouflage: Photonic crystals in the diamond weevil, *Entimus imperialis*. Proceedings of the Royal Society B: Biological Sciences 279(1738): 2524–2530. <https://doi.org/10.1098/rspb.2011.2651>
- Zhang G, Basharat U, Matzke N, Franz NM (2017) Model selection in statistical historical biogeography of Neotropical insects–The *Exophthalmus* genus complex (Curculionidae: Entiminae). Molecular Phylogenetics and Evolution 109: 226–239. <https://doi.org/10.1016/j.ympev.2016.12.039>

Supplementary material 1

DarwinCore-formatted list of species of Entiminae recorded from Colombia. This checklist is available via GBIF (Girón 2020, <https://doi.org/10.15472/jdwfao>)

Authors: Jennifer C. Girón

Data type: Species list

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/neotropical.15.59713.suppl1>

Supplementary material 2

DarwinCore-formatted occurrence records for 749 entomine specimens deposited in Colombian biological collections

Authors: Jennifer C. Girón

Data type: Occurrence data

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/neotropical.15.59713.suppl2>